

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problems Mailbox.**



Pergamon

0163-7258(94)00048-4

Pharmac. Ther. Vol. 64, pp. 445-475, 1994  
Copyright © 1994 Elsevier Science Ltd  
Printed in Great Britain. All rights reserved  
0163-7258/94 \$26.00

Specialist Subject Editor: F. MITCHELSON

## PURINOCEPTORS: ARE THERE FAMILIES OF P2X AND P2Y PURINOCEPTORS?

MARIA P. ABBRACCHIO\* and GEOFFREY BURNSTOCK†

\*Institute of Pharmacological Sciences, University of Milan, Via Balzaretto 9,  
20133 Milan, Italy

†Department of Anatomy and Developmental Biology, University College London,  
Gower Street, London WC1E 6BT, U.K.

**Abstract**—There has been an exponential growth in interest in purinoceptors since the potent effects of purines were first reported in 1929 and purinoceptors defined in 1978. A distinction between P<sub>1</sub> (adenosine) and P<sub>2</sub> (ATP/ADP) purinoceptors was recognized at that time and later, A<sub>1</sub> and A<sub>2</sub> as well as P<sub>1x</sub> and P<sub>1y</sub> subclasses of P<sub>1</sub> and P<sub>2</sub> purinoceptors were also defined. However, in recent years, many new subclasses have been claimed, particularly for the receptors to nucleotides, including P<sub>2a</sub>, P<sub>2b</sub>, P<sub>2c</sub> and P<sub>2d</sub>, and there is some confusion now about how to incorporate additional discoveries concerning the responses of different tissues to purines. The studies beginning to appear defining the molecular structure of P<sub>2</sub>-purinoceptor subtypes are clearly going to be important in resolving this problem, as well as the introduction of new compounds that can discriminate pharmacologically between subtypes. Thus, in this review, on the basis of this new data and after a detailed analysis of the literature, we propose that:

- (1) P2X(ligand-gated) and P2Y(G-protein-coupled) purinoceptor families are established;
- (2) four subclasses of P2X-purinoceptor can be identified (P2X<sub>1</sub>, P2X<sub>2</sub>) to date;
- (3) the variously named P<sub>2</sub>-purinoceptors that are G-protein-coupled should be incorporated into numbered subclasses of the P2Y family. Thus:

P2Y<sub>1</sub> represents the recently cloned P2Y receptor (clone 803) from chick brain;  
P2Y<sub>2</sub> represents the recently cloned P<sub>2a</sub> (or P<sub>2c</sub>) receptor from neuroblastoma, human  
epithelial and rat heart cells;

P2Y<sub>3</sub> represents the recently cloned P2Y receptor (clone 103) from chick brain that resembles  
the former P<sub>2a</sub> receptor;

P2Y<sub>4</sub>, P2Y<sub>5</sub> represent subclasses based on agonist potencies of newly synthesised analogues;  
P2Y<sub>6</sub> represents the 10mer P<sub>2d</sub> receptor for dinucleotides.

This new framework for P<sub>2</sub> purinoceptors would be fully consistent with what is emerging for the receptors to other major transmitters, such as acetylcholine,  $\gamma$ -aminobutyric acid, glutamate and serotonin, where two main receptor families have been recognised, one mediating fast receptor responses directly linked to an ion channel, the other mediating slower responses through G-proteins. We fully expect discussion on the numbering of the different receptor subtypes within the P2X and P2Y families, but believe that this new way of defining receptors for nucleotides, based on agonist potency order, transduction mechanisms and molecular structure, will give a more ordered and logical approach to accommodating new findings. Moreover, based on the extensive literature analysis that led to this proposal, we suggest that the development of selective antagonists for the different P<sub>2</sub>-purinoceptor subtypes is now highly desirable, particularly for therapeutic purposes.

**Keywords**—ATP, P<sub>2</sub> purinoceptors, P2X and P2Y, ligand-gated channels, G-protein-linked receptors.

**Abbreviations**— $\alpha$ -GMP,  $\alpha$ -glycylglycine; ATP, ADP, adenosine 5'-O-(2-thiodiphosphate); ANAPP<sub>3</sub>, 3-O-[N-(4-azido-2-nitrophenyl)amino] propionyl ATP; AppNHp, adenosine 5'[( $\beta$ , $\gamma$ -imido)triphosphate]; ApxA, diadenosine polyphosphate; ATP<sub>2</sub>S, adenosine 5'-O-(1-thiotriphosphate); ATP<sub>3</sub>S, adenosine 5'-O-(3-thiotriphosphate);  $\beta$ -GMP,  $\beta$ -glycylglycine; CF, cystic fibrosis; DAG, diacylglycerol; 2mSATP, 2-methylthio-ATP; IP<sub>3</sub>, inositol-1,4,5-trisphosphate; PPADS, pyridoxalphosphate-6-azophenyl-2',4'-disulfonic acid.

## CONTENTS

1. Early History of Receptors to Nucleosides and Nucleotides	446
1.1. $P_1$ and $P_2$ purinoceptors	446
1.2. Subclasses of the adenosine/ $P_1$ purinoceptor	447
1.3. Subclasses of the ATP/ $P_2$ purinoceptor	447
2. Molecular Biology of ATP Purinoceptors	450
3. New ATP Derivatives as Possible Tools to Discriminate $P_2$ Purinoceptor Subtypes	455
4. New Proposals for $P_2$ -Purinoceptor Classification	458
4.1. $P_2$ -purinoceptor coupling to different transduction mechanisms: a criterion for distinguishing the $P_2X$ -(ligand-gated) and $P_2Y$ -(G-protein-coupled) purinoceptor families	458
4.2. Do the $P_{2a}$ , $P_{2b}$ and $P_{2c}$ receptors belong to the $P_2Y$ -purinoceptor family?	458
4.3. Proposals for the subclassification of the $P_2Y$ -purinoceptor family	459
4.4. Proposals for the subclassification of the $P_2X$ -purinoceptor family	463
5. $P_2$ -Purinoceptor Ligands as Potential Therapeutic Entities	465
5.1. Diabetes	465
5.2. Cancer	466
5.3. Cystic fibrosis	466
5.4. Pulmonary hypertension	466
5.5. Surfactant and mucin secretion	466
5.6. Renal failure	467
5.7. Bone and cartilage diseases	467
5.8. Urinary incontinence	467
5.9. Thrombosis	467
5.10. Ventricular tachycardia	467
5.11. Gastro-intestinal tract	468
5.12. Centrally targeted $P_2$ purinoceptor drugs	468
5.13. Human reproduction	468
6. Future Developments	468
Note Added in Proof	468
Acknowledgments	468
References	468

## 1. EARLY HISTORY OF RECEPTORS TO NUCLEOSIDES AND NUCLEOTIDES

1.1.  $P_1$  and  $P_2$  Purinoceptors

The potent extracellular actions of purine nucleotides and nucleosides, namely adenosine and ATP, were first recognised by Drury and Szent-Gyorgyi (1929) in a seminal paper reporting the activities of adenine compounds on the mammalian heart. Following this report, there was considerable activity within the field, with particular emphasis on the actions of adenosine and ATP on the cardiovascular system (Gillespie, 1933; Green and Stoner, 1930).

The potent vasodilator actions of adenylyl compounds led Holton and Holton (1954) to suggest that ATP might represent the vasodilatory substance that was released upon antidromic stimulation of sensory nerves supplying the rabbit ear artery, a hypothesis that was later demonstrated following stimulation of the great auricular nerve (Holton, 1959). The sensitivity of the coronary vasculature to these same compounds prompted Berne (1963) to propose that adenosine was the physiological mediator of the coronary vasodilation associated with myocardial hypoxia, which further supported the role of ATP and adenosine in physiological regulatory processes.

In the same period (early 1960s), a component that was neither adrenergic nor cholinergic was recognised in the autonomic nervous system. This component was strongly represented in the gastrointestinal tract and later identified in a variety of organs, including lung, bladder, seminal vesicles, esophagus, uterus, eye, trachea and part of the cardiovascular system. Using the criteria summarised by Eccles (1964) for the acceptance of putative neurotransmitters, in the early 1970s, Burnstock proposed that the principal active substance released from at least some of these nerves was ATP and, as a consequence, these nonadrenergic, noncholinergic nerves were tentatively termed 'purinergic' (Burnstock *et al.*, 1970; Burnstock, 1972).

In the subsequent years, the purinergic theory was strengthened by a great body of experimental evidence supporting the role of ATP as a transmitter or co-transmitter with noradrenaline,

Purinoceptors: Are there families of  $P_2X$  and  $P_2Y$  purinoceptors?

447

acetylcholine and other substances (Burnstock, 1976, 1990) and by the identification of specific extracellular receptors mediating the variety of physiological effects induced by purines.

Based largely on an analysis of the voluminous literature about the actions of purines on a wide number of tissues, in 1978, in a seminal review, Burnstock proposed a basis for distinguishing two types of purinergic receptors, termed  $P_1$  and  $P_2$ , the purinoceptor preferentially activated by adenosine and ATP, respectively. The original classification into  $P_1$  and  $P_2$  purinoceptors was based on four criteria: (1) the relative potencies of ATP, ADP, AMP and adenosine; (2) the selective actions of antagonists, particularly methylxanthines, which competitively antagonise adenosine, but not ATP actions; (3) the modulation of adenylate cyclase with resultant changes in intracellular cAMP levels by adenosine, but not ATP, and (4) the induction of prostaglandin synthesis by ATP, but not by adenosine. Since the time of this proposal, many experiments have been carried out that support and extend the  $P_1/P_2$  classification (Williams, 1987; Burnstock, 1991), which is now well-established and largely used in the literature and has been adopted by the IUPHAR Subcommittee for Purinoceptor Subclassification (Abbraccio *et al.*, 1993; Fredholm *et al.*, 1994).

In general, the prejunctional purinoceptors that modulate release of noradrenaline from postganglionic sympathetic nerves are  $P_1$ . Westfall *et al.* (1990) have proposed the existence of a third purinoceptor subtype on some sympathetic nerve terminals tentatively named  $P_3$  and claimed to recognise the structure of both nucleotides and nucleosides; however, this proposal has not received wide acceptance as yet.

1.2. Subclasses of the Adenosine/ $P_1$  Purinoceptor

Following Burnstock's proposal, subsequent work led to the demonstration of both adenosine/ $P_1$  and ATP/ $P_2$  purinoceptor subtypes.

Parallel work by Lontos *et al.* (1980) and Hamprecht's group (van Calker *et al.*, 1979) resulted in the identification of two subclasses of adenosine receptors, which were originally defined on the basis of whether their activation inhibited or stimulated adenylate cyclase activity. Lontos *et al.* termed these receptors  $R_i$  and  $R_s$ , based on the need of an intact ribose ring in the purine, with the subscripts 'i' and 's' referring to inhibition and activation of cAMP formation, respectively, whereas these same two receptors were named  $A_1$  and  $A_2$  by van Calker *et al.*, a nomenclature that found preference in the subsequent literature (Abbraccio *et al.*, 1993).

More recently,  $A_1$  and  $A_2$  receptor subtypes have been identified by cloning (Zhou *et al.*, 1992) and binding studies (Cornfield *et al.*, 1992), respectively. A detailed analysis of experimental evidence supporting the subclassification of the  $P_1$  purinoceptor and of the pathophysiological roles of the different receptor subtypes is not within the aims of the present review, and the reader is referred to excellent reviews recently published in this field (Jacobson *et al.*, 1992; Linden *et al.*, 1994).

1.3. Subclasses of the ATP/ $P_2$  Purinoceptor

Burnstock and Kennedy (1985) provided evidence for a subclassification of the  $P_2$  receptor into the  $P_{2a}$  and  $P_{2b}$  subtypes. In this first in-depth review of the functional effects of various ATP analogues in a number of different biological systems, they discriminated between two major classes of receptor-mediated responses on the basis of a different response profile to ATP analogs and selective antagonism. Thus, for  $P_{2a}$  purinoceptors, ATP analogs may be listed in order of potency as follows:  $\alpha\beta\text{meATP} \geq \beta\text{meATP} > \text{ATP} \approx 2\text{meSATP} = \text{ADP}$ ;  $3\text{-O-}\beta\text{N-(4-azido-2-nitro-phenyl)amino} \text{ propionyl ATP (ANAPP3)}$  was claimed to behave as a selective antagonist, whereas prolonged exposure to  $\alpha\beta\text{-methylene ATP (}\alpha\beta\text{meATP)}$  selectively desensitises this receptor (Kasakov and Burnstock, 1983). For  $P_{2b}$  purinoceptors,  $2\text{meSATP} > \text{ATP} > \alpha\beta\text{meATP} = \beta\text{meATP}$ ; reactive blue 2, an anthraquinone sulfonic acid derivative, has been claimed to be a selective antagonist, at least over a limited concentration range (Kerr and Krantis, 1979; Manzini *et al.*, 1986; Houston *et al.*, 1987).

Studies of the pharmacological actions of isopolar phosphonate analogues of ATP in guinea-pig

taenia coli and bladder have supported the  $P_{2X}/P_{2Y}$  subclassification of  $P_2$  purinoceptors in smooth muscle and have also shown that  $\beta$ -methylene ATP ( $\beta$ meATP) and its analogues behave as selective  $P_{2X}$  agonists (Cusack *et al.*, 1987), while adenosine 5'-O-(2-thiodiphosphate) (ADP $\beta$ S) is a specific agonist at  $P_{2Y}$ -purinoceptors (Hourani *et al.*, 1988). More recently, the trypanoside suramin has been shown to be a specific antagonist for  $P_2$ -purinoceptors, although not selective for the  $P_{2X}$  and  $P_{2Y}$  subclasses (Hoyle *et al.*, 1990), whereas the new compound pyridoxalphosphate-6-azophenyl-2',4'-disulfonic acid (PPADS) synthesised by Lambrecht's group appears to represent a selective  $P_{2X}$  antagonist (Lambrecht *et al.*, 1992; Ziganshin *et al.*, 1993, 1994b).

After Burnstock and Kennedy's proposal, in 1986, Gordon further delineated ATP effects in a variety of tissues (Gordon, 1986) and defined two additional  $P_2$  purinoceptor subtypes on platelets and mast cells (and lymphocytes) that did not seem to fit the  $P_{2X}/P_{2Y}$  subclassification and were tentatively termed  $P_3$  and  $P_4$  purinoceptors, respectively. The platelet receptor is unique, being activated by ADP and blocked by ATP, whereas the mast cell and macrophage ATP receptor seems to be preferentially activated by the tetrabasic form of ATP, ATP $^{4-}$ , and irreversibly inhibited by oxidised ATP (Murgia *et al.*, 1993). Doubts have been raised about the actual 'receptor' nature of the  $P_3$  purinoceptor subtype, since it is represented by the opening of a fairly nonselective membrane pore. Some of the characteristics of the different  $P_2$  purinoceptor subtypes are summarised in Table 1.

From studies performed after Burnstock and Kennedy's proposal, it was evident that the  $P_{2X}$  and  $P_{2Y}$  purinoceptors could also be differentiated on the basis of their transduction mechanisms. The  $P_{2X}$  purinoceptor subtype involves an intrinsic ion channel permeable to Na $^{+}$ , K $^{+}$  and Ca $^{2+}$  (Benham and Tsien, 1987; Bean, 1992), whereas the  $P_{2Y}$  purinoceptor subtype is a G-protein coupled receptor that modulates membrane phosphoinositide metabolism and, hence, inositol-1,4,5-trisphosphate (IP3) and diacyl-glycerol (DAG) generation (Cusack, 1993), although additional transduction mechanisms also seem to be utilised in some tissues, such as modulation of cAMP generation (Okajima *et al.*, 1989; Yamada *et al.*, 1992) and arachidonic acid mobilisation (Brunner and Murphy, 1990).

There is now good evidence in favour of the existence of purinoceptors that respond to the pyrimidine derivative UTP, as well as to ATP and adenosine 5'-O-(3-thiotriphosphate) (ATP $\gamma$ S) (but not to ZmeSA TP or  $\alpha$ meATP, Table 1). As for the  $P_{2Y}$  and  $P_{2X}$  subtypes, this receptor is linked to G-protein activation (Table 1) and has been alternatively termed ' $P_{2Y}$ ' or 'nucleotide' or 'pyrimidine' receptor. At this receptor subtype, UTP is generally either equipotent or more potent than ATP, suggesting that the 'pyrimidine/nucleotide' receptor might indeed represent a further subclass of the  $P_2$  purinoceptor (see also Section 2).

Besides these  $P_2$  purinoceptor subtypes, there also appear to be receptors for adenine dinucleotide polyphosphates (Ap4A, Ap5A and Ap6A), leading to the proposal of an additional receptor subtype, the ' $P_{2D}$ ' purinoceptor (Hilderman *et al.*, 1991; Castro *et al.*, 1992; Table 1). The physiological functions of the diadenosine polyphosphate (ApxA) compounds are unclear at present, although possible roles as neurotransmitters, co-transmitters or trophic factors have been suggested (Hoyle, 1990).

The use of different terminologies to define UTP-sensitive responses and the apparent random walk through the alphabet in the naming of  $P_{2X}$ ,  $P_{2Y}$ ,  $P_{2D}$  and  $P_{2Z}$  and  $P_{2B}$  has generated some concern in the IUPHAR Subcommittee for Purinoceptor Subclassification (Abbracchio *et al.*, 1993; Fredholm *et al.*, 1994), and has led to confusion and inconsistency, especially when trying to incorporate into current nomenclature schemes additional discoveries about the biological responses to purines and pyrimidines.

This confusion has prompted us to perform a detailed analysis of the available literature in order to verify whether a new way of defining the different receptor subtypes for nucleotides could be proposed. Knowledge of receptor molecular structure is critical and strategic in defining neurotransmitter receptor subtypes. Since molecular biology data on  $P_2$  purinoceptors are now becoming available, we would like to summarise initially such available information in order to try to identify some useful criteria in  $P_2$  purinoceptor subclassification. Further, we report here the results of studies recently performed with a number of modified ATP analogues that give further hints about possible  $P_{2X}$ - and  $P_{2Y}$ -purinoceptor subclasses.

Table 1.  $P_2$  Purinoceptor: Characteristics and Current Subclassification

Receptor	$P_{2X}$	$P_{2Y}$	$P_{2B}$ ( $P_{2D}$ )	$P_{2Z}$	$P_{2L}$	$P_{2D}$
Type	Intrinsic ion channel (Na $^{+}$ , K $^{+}$ , Ca $^{2+}$ )	G-protein-coupled (IP3/Ca $^{2+}$ /DAG)	G-protein-coupled (IP3/Ca $^{2+}$ /DAG)	G-protein-coupled (IP3/Ca $^{2+}$ /DAG/cAMP)	Nonselective pore	G-protein-coupled (IP3/Ca $^{2+}$ /DAG)
Agonist profile	$\alpha$ meATP $\geq$ $\beta$ meATP $>$ ATP $\geq$ 2meSATP $\approx$ ADP	2meSATP $\gg$ ATP $\gg$ $\alpha$ meATP	UTP $\geq$ ATP $\gg$ 2meSATP	2-ADP	ATP $^{4-}$	ApxA
Antagonist	Desensitisation by $\alpha$ meATP; blocked by suramin, by ANAPP3 and PPADS	Blocked by suramin and by reactive blue 2		ATP	Oxidised ATP	






Fig. 1(A, B)—see opposite.

## 2. MOLECULAR BIOLOGY OF ATP PURINOCEPTORS

Acquired ATP-generated currents in *Xenopus* oocytes injected with mRNA from embryonic guinea-pig brain were first demonstrated by Fournier *et al.* (1990), and by Honoré *et al.* (1991), and also following injection of size-fractionated RNA from J774 murine macrophage-like cells by Hickman *et al.* (1993). ATP<sub>2</sub>S-induced acceleration of intracellular calcium efflux was shown in *Xenopus* oocyte injected with mRNA from HL60 promyelocytic leukaemia cells, a response that is believed to be mediated by a G-protein-linked P<sub>2</sub> purinoceptor (Murphy and Tiffany, 1990). More recently, Russell *et al.* (1993) reported ATP-generated currents in *Xenopus* oocytes injected with poly-(A)<sup>+</sup> mRNA from guinea-pig vas deferens. Such ATP-generated currents involved the release of intracellular Ca<sup>2+</sup> and also depended upon Ca<sup>2+</sup> influx.

The Barnard and Burnstock groups have recently cloned a P<sub>2</sub>-like purinoceptor using a homology screening strategy. A previously isolated guinea-pig partial cDNA,\* the sequence homologue

\*Webb, T. E., Bateson, A. N. and Barnard, E. A. (1991) A polymerase chain reaction based strategy for the isolation of DNA sequences that encode G-protein-coupled receptors. In: *17th EMBO Annual Symposium, Molecular Mechanisms of Signal Transduction*, Heidelberg, Germany, 16–19 September, 1991, Abstract No. 208.




Fig. 1(C). Responses in cRNA-injected *Xenopus* oocytes (at  $-40$  mV). Aa: Dose-dependence of membrane currents evoked by ATP ( $10$ – $300$   $\mu$ M). The transient downward deflections monitor the input conductance following hyperpolarising voltage steps ( $-10$  mV, applied every 5 sec for 1 sec). Similar results were observed in two other oocytes. Ab, Ac: In two oocytes, the agonist selectivity was assessed using 2MeS-ATP, ATP, ADP,  $\alpha$ Me-ATP and UTP. Holding potentials,  $-40$  mV. B: ATP concentration–response relationship. The membrane current amplitude ( $I_{ATP}$ ) was measured using the initial peak inward current induced by each ATP concentration. C: Suramin antagonised the responses to (Ca) ATP and (Cb) 2MeS-ATP. Reactive blue 2 (RB) also inhibited the responses to (Cc) ATP and (Cd) 2MeS-ATP. Bar indicates ligand application. Adapted from Webb *et al.* (1993).

of the canine orphan receptor RDC1 (Libert *et al.*, 1989), was used to screen an embryonic chick whole brain cDNA library under conditions of low stringency (Webb *et al.*, 1993). The cloning strategy included use of two degenerate oligonucleotide primers from transmembrane domains II and VI of G-protein-coupled receptors for polymerase chain reaction amplification of guinea-pig brain first-strand cDNA, subcloning of amplification products and screening of an embryonic chick whole brain cDNA library.

An isolated clone, 803, encoded a 362-amino acid polypeptide displaying the typical topology of a G-protein-coupled receptor (Webb *et al.*, 1993), such as the presence of seven hydrophobic putative transmembrane  $\alpha$ -helices; consensus sequences for N-linked glycosylation in the N-terminus extracellular domain; two conserved cysteine residues in the first and second extracellular loops, believed to form disulphide bonds; and several consensus phosphorylation sites in the third cytoplasmic loop and C-terminal intracellular domain (Linden *et al.*, 1991).

Functional expression of this clone in *Xenopus laevis* indicated that the encoded protein is a P<sub>2</sub>-purinoceptor. ATP induced a large calcium-activated inward chloride current (Webb *et al.*, 1993; Fig. 1Aa). Responses to ATP were dose-dependent with an EC<sub>50</sub> of  $49.5 \pm 6$   $\mu$ M (Fig. 1Aa, B). Further analysis of the ligand specificity of this receptor using a range of P<sub>2</sub>-purinoceptor agonists led to designation of this clone as a P<sub>2</sub>Y-like purinoceptor. ATP-like effects were also induced by 2MeS-ATP

and ADP, whereas both UTP and the P<sub>x</sub> selective agonists  $\alpha$ MeATP and  $\beta$ MeATP were inactive at concentrations up to 30–100  $\mu$ M. Therefore, the rank order of potency of these agonists was: 2MeSATP  $\geq$  ATP  $>$  ADP  $\gg$   $\alpha$ MeATP,  $\beta$ MeATP, UTP (Fig. 1A,B,C). Both suramin and the P<sub>1</sub>-selective antagonist reactive blue 2 antagonised the responses to ATP and 2MeSATP (Fig. 1C). Based on selective antagonism by reactive blue 2, high potency of ATP, inactivity of UTP and methylene ATP derivatives, and strong structure similarity with G-protein-coupled receptors, the newly cloned receptor was identified as a P<sub>2</sub> subtype rather than a P<sub>x</sub> or P<sub>u</sub>/nucleotide purinoceptor subtype. Moreover, the near equipotency of 2MeSATP and ATP and greater potency of ATP over ADP suggested that this expressed receptor was a novel subtype of the P<sub>2</sub>Y purinoceptor family, which was consequently named 'P<sub>2</sub>Y<sub>1</sub>'.

The agonist profile typical of this receptor (2MeSATP  $\geq$  ATP  $>$  ADP) was a starting point for us in analysing the available literature in our effort to find a new and logical way of defining the different receptor subtypes to ATP (see Section 4.4).

The cloning work by the Barnard and Burnstock groups also led to the isolation of an ADP-sensitive clone (clone 103), which, based on preliminary pharmacological results, might be related to the so-called 'P<sub>2</sub>' purinoceptor (Barnard *et al.*, 1994).

Lustig *et al.* (1993) have independently cloned a cDNA from NG 108-15 neuroblastoma cells encoding a metabotropic P<sub>2</sub>/nucleotide purinoceptor by utilising a *X. laevis* oocyte expression cloning strategy. In *Xenopus* oocytes injected with poly(A)<sup>+</sup> RNA from NG 108-15 cells, ATP or UTP (1 mM) evoked calcium-dependent inward currents. To identify the cDNA encoding the receptor protein responsible for these currents, pools of  $2 \times 10^5$  individual clones from an NG 108-15 cDNA plasmid library were *in vitro* transcribed and the resultant cRNA was injected into oocytes. The pool that rendered the oocytes responsive to ATP or UTP was subdivided progressively and the procedure was repeated until a single clone, pP2R, was obtained (Lustig *et al.*, 1993).

In oocytes injected with pP2R cRNA transcripts, inward currents were elicited by bath application of ATP, UTP or ATP<sub>S</sub>, but not by comparable concentrations of 2MeSATP, ADP,  $\beta$ MeATP or  $\alpha$ MeATP. The order of agonist potency was ATP  $\approx$  UTP  $>$  ATP<sub>S</sub>  $\gg$  2MeSATP, ADP,  $\beta$ MeATP,  $\alpha$ MeATP, which, again, was a starting point for us in analysing the effects reported in the literature for ATP and UTP.

Recently, the ATP/UTP receptor has been cloned from human airway and colonic epithelium (Parr *et al.*, 1994) and from rat heart (Godecke and Schrader, 1994). Similarly to the receptor cloned by Lustig *et al.*, the HP2U clone isolated by Parr *et al.*, when expressed in 1321N1 astrocytoma cells, displayed comparable responses to ATP and UTP, whereas 2MeSATP and  $\alpha$ MeATP had little effect. Activation of this receptor leads to breakdown of phosphoinositides and increase of intracellular Ca<sup>2+</sup>, an effect that was partially blocked by pertussis toxin.

Both the P<sub>2</sub>Y<sub>1</sub> purinoceptor and the cloned pP2R and HP2U purinoceptors showed predicted structural features characteristic of most known G-protein-coupled receptors.

Figure 2 shows the deduced protein sequences of cloned P<sub>2</sub>Y<sub>1</sub> and pP2R purinoceptors in comparison with adenosine A<sub>1</sub>, A<sub>2</sub>, A<sub>3</sub> and A<sub>4</sub> receptors. Interestingly, the P<sub>2</sub>Y<sub>1</sub> and pP2R sequences differ markedly from the cloned adenosine receptors. In fact, the pP2R receptor is more similar to receptors for various peptides, such as thrombin (25% identity), platelet-activating factor (25% identity), angiotensin II (22% identity), interleukin 8 (23% identity) and GPRN1, a putative vasoreactive intestinal peptide receptor (21% identity) (Lustig *et al.*, 1993). The cloned receptor is substantially less similar (<12% identity) to adenosine and cAMP receptors.

Similarly, the P<sub>2</sub>Y<sub>1</sub> receptor has only a low sequence identity with the adenosine A<sub>1</sub> (21%) and the cAMP (17%) receptors, whereas, as reported for the pP2R receptor, higher identities were found with the angiotensin II, thrombin, platelet-activating factor and interleukin 8 receptors (Table 2), suggesting that these two purinoceptors might belong to a completely different subfamily of the G-protein-coupled receptor class.

The tissue distribution of the P<sub>2</sub>Y<sub>1</sub> transcript determined by Northern hybridisation revealed a discrete pattern of expression in the adult chicken. The P<sub>2</sub>Y<sub>1</sub> mRNA was present in brain, spinal cord, gastrointestinal tract, spleen and leg muscle (Webb *et al.*, 1993). Conversely, for the pP2R receptor, Northern blot analysis revealed wide distribution of mRNA in all tested mouse tissues, such as spleen, testes, kidney, liver, lung, heart and brain (Lustig *et al.*, 1993). This distribution is very similar to that reported for the HP2U receptor (Parr *et al.*, 1994). The distribution of the P<sub>2</sub>Y<sub>1</sub> (Webb

TM 1		TM 2	
A	A1-RECEPTORS	MPPSISAFQAAIIGIEVLIALVSPGVNLVIAVAVKVNQALRDATFCFIVSLAVADVAVGALVIFLAILINIGPQTYFHTC	
	hA1	MPPYISAFQAAIIGIEVLIALVSPGVNLVIAVAVKVNQALRDATFCFIVSLAVADVAVGALVIFLAILINIGPQTYFHTC	
	rA1	MPPAISAFQAAIIGIEVLIALVSPGVNLVIAVAVKVNQALRDATFCFIVSLAVADVAVGALVIFLAILINIGPRTYFHTC	
	dA1	MPPSISAFQAAIIGIEVLIALVSPGVNLVIAVAVKVNQALRDATFCFIVSLAVADVAVGALVIFLAILINIGPRTYFHTC	
	A2-RECEPTORS	MPINGSSVYITVELAIAVLAILQNVLVCAVWVLSNQLQNVNIFVVSLLAAADIAVGVLAIPTAITISTGFCACCHGC	
	hA2a	MSTNGSSVYITVELAIAVLAILQNVLVCAVWVLSNQLQNVNIFVVSLLAAADIAVGVLAIPTAITISTGFCACCHNC	
	rA2a	MGSSVYITVELAIAVLAILQNVLVCAVWVLSNQLQNVNIFVVSLLAAADIAVGVLAIPTAITISTGFCACCHGC	
	hA2b	MLLETQDALYVALELVIAALAVAGNVLVCAVAGTANTLTPTNYFLVSLAAADVAVGLFAIPTAITISLGFCDFYGC	
	A3-RECEPTORS	MQLETQDALYVALELVIAALAVAGNVLVCAVAGTANTLTPTNYFLVSLATADVAVGLFAIPTAITISLGFCDFHSC	
	rA3	MKANNTTTSALNLOITTYTNEAAIGLCAVVGQMLVIMVVKLNRLTRTTTFYFIVSLALADIAVGVLVIFLAIASVLEVMHFYAC	
	P2-RECEPTORS	MTEALISAALNGTQPELLAGGMAAGNATTECSLTNTGTFQFYLLPTVYILVFTITGFLGHSVAIMNFVFMHRPWSQISVDMNLALADFLVLTLPALIFYYFNKTDNIFGDMCK	
	P2Y1	MAADLEPNWSTINGTWEGDELGYKCRFNEDEXTVLLPVSYGVVCLGLCLNVVLYIFLCRLKT-WNASTTMMFLAVSDSLYASLPLLVYVYARGDHPFSTVLCK	
	P2U	TM 3	
		TM 4	
	hA1	LMVACPVLILTQSSILALLAIAVDRLRVKIPRLRYKVVTPPRAAVAIAGCWLSFVGLTPTGWN.....NLSAVERAANAANGSGEPVICKFEFENVISMEYMYINFTFWVLPP	
	rA1	LMVACPVLILTQSSILALLAIAVDRLRVKIPRLRYKVVTPPRAAVAIAGCWLSFVGLTPTGWN.....NLSVVEQDWRANGCVGEPVICKFEFENVISMEYMYINFTFWVLPP	
	dA1	LMVACPVLILTQSSILALLAIAVDRLRVKIPRLRYKVVTPPRAAVAIAGCWLSFVGLTPTGWN.....RLGEAQRANAANGSGEPVICKFEFENVISMEYMYINFTFWVLPP	
	hA1	LMVACPVLILTQSSILALLAIAVDRLRVKIPRLRYKVVTPPRAAVAIAGCWLSFVGLTPTGWN.....NLSAVERDNLANGSVGEPVIECQFEKVISMEYMYINFTFWVLPP	
	hA2a	LFIACFVLVLTQSSIFSLAIAIDRYIAIRIPRLRYNGLVTGTRAGGIIAICWLSFAIGLTPFLGWN.....NCSQPKQEGKNHSQCGGEGQVACLFEDVVPMMYINFTFWVLPP	
	rA2a	LFIACFVLVLTQSSIFSLAIAIDRYIAIRIPRLRYNGLVTGTRAGGIIAICWLSFAIGLTPFLGWN.....NCSQPKQEGKNHSQCGGEGQVACLFEDVVPMMYINFTFWVLPP	
	rA2a	LFIACFVLVLTQSSIFSLAIAIDRYIAIRIPRLRYNGLVTGTRAGGIIAICWLSFAIGLTPFLGWN.....NCSQKDGNSKT..CGEGRVTCLEFEDVVPMMYINFTFWVLPP	
	hA2b	LFIACFVLVLTQSSIFSLAIAVDRLAICVFLRYKSLVTGTRAGGIIAICWLSFAIGLTPFLGWN..DSATNNCTEPWDGTTNE..SCC..LVKCLFENVVPMYSIMYINFTFWVLPP	
	rA2b	LFIACFVLVLTQSSIFSLAIAVDRLAICVFLRYKSLVTGTRAGGIIAICWLSFAIGLTPFLGWN..DRATSNCTEPGDGTTNK..SCC..PVKCLFENVVPMYSIMYINFTFWVLPP	
	rA2b	LFIACFVLVLTQSSIFSLAIAVDRLAICVFLRYKSLVTGTRAGGIIAICWLSFAIGLTPFLGWN..DRATSNCTEPGDGTTNK..SCC..PVKCLFENVVPMYSIMYINFTFWVLPP	
	rA3	LFMSCVLLVTFTHASIMSLAIAVDRLRVKLTVRVYRTVTTQRRWLFGLGCLVSLVGLTPTGWNRRKVTLELSONSS..TLS.....CHFRFVGLDXMVTTSFTTWILIP	
	P2Y1	LQRFIFHVNLVGSILFLTCISVHRYTGV...VHPLKSLGRLKKNAVYVSLVVALVAVIADIFYSGTGVRR..NKT....ITCYDTTADDEY...LRSTFVYSNCTVTHFCIPP	
	P2U	LVRFIFVNLVGSILFLTCISVHRCGLV..RPLHSLRWGRA..RYARRVAVVVVLVLAQAPVLYFVTTSVRGTRITCHDTSARELFS	
		HIVAYSSVKGLLI	
		TM 5	

Fig. 2(A).

	TM 5	TM 6	TM 7
hA1	LLIMVLIYLEVYFLIRKQLNKKVSA..SSGDPQKYGGKELKIAKSLALILFLALSWLPLHILNCITLFCPSC..HKPSILTYIAIFLTHGSAAGPIVIAF		
rA1	LLIMVLIYLEVYFLIRKQLNKKVSA..SSGDPQKYGGKELKIAKSLALILFLALSWLPLHILNCITLFCPTC..QKPSILTYIAIFLTHGSAAGPIVIAF		
dA1	LLIMVLIYLEVYFLIRKQLGKKVSA..SSGDPQKYGGKELKIAKSLALILFLALSWLPLHILNCITLFCPSC..RKPSILTYIAIFLTHGSAAGPIVIAF		
bA1	LLIMVLIYMEVYFLIRKQLSKKVSA..SSGDPQKYGGKELKIAKSLALILFLALSWLPLHILNCITLFCPSC..HMPRIIYIAIFLTHGSAAGPIVIAF		
hA2a	LLIMLVYVLRIFLAARRQLKQMESQPLPGERARSTLQKEVHAAKSLAIVGLFALCWLPLHILNCITLFCPSC..HAPLWIMYLAIVLSHNSVVMPIVYAY		
dA2a	LLIMLVYVLRIFLAARRQLKQMESQPLPGERARSTLQKEVHAAKSLAIVGLFALCWLPLHILNCITLFCPSC..HAPLWIMYLAIVLSHNSVVMPIVYAY		
rA2a	LLIMLVYVLRIFLAARRQLKQMESQPLPGERARSTLQKEVHAAKSLAIVGLFALCWLPLHILNCITLFCPSC..HAPLWIMYLAIVLSHNSVVMPIVYAY		
hA2b	LLIMLVYVLRIFLAARRQLKQMESQPLPGERARSTLQKEVHAAKSLAIVGLFALCWLPLHILNCITLFCPSC..HAPLWIMYLAIVLSHNSVVMPIVYAY		
rA2b	LLIMLVYVLRIFLAARRQLKQMESQPLPGERARSTLQKEVHAAKSLAIVGLFALCWLPLHILNCITLFCPSC..HAPLWIMYLAIVLSHNSVVMPIVYAY		
rA3	LVMNCIITLDIFYIIRNKLSQNLTG...FRETRAIFYGREFKTAKSLFLVLFLALCWLPLHILNCITLFCPSC..KIPRIAMCLOILLSHNSVVMPIVYAY		
P2Y1	IVILQCYGLIVKALIKDLNDSF.....LRRKSIYLVIIIVLTAVSYLPLHILNCITLFCPSC..DFQTPQMCANDKVYATIQVTRGLASLHNSVVMPIVYAY		
P2U	FAVPSVILVCTVIMARRLLKPAYGTTGGLPRAKRA.....SVRTIALVIAVFLCFLPLHILNCITLFCPSC..DFQTPQMCANDKVYATIQVTRGLASLHNSVVMPIVYAY		
hA1	RIQKFRVTFKLIWNDHFRCPAPPIDEDLFEERPD		
rA1	RIHKKFRVTFKLIWNDHFRCPAPPIDEDLFEERPD		
dA1	RIQKFRVTFKLIWNDHFRCPAPPIDEDLFEERPD		
bA1	RIQKFRVTFKLIWNDHFRCPAPPIDEDLFEERPD		
hA2a	RIREFRQTFRKIIIRSHVLRQEPFKAAGTSARVLAAGSDGEQVSLRLNGHPGCVWANGSAPHERRPNGYALGLVSGGSAQESQGNLGLPDVELLSHELKGVCEPPGLDDPLAQDGAGVS		
dA2a	RIREFRQTFRKIIIRSHVLRQEPFKAAGTSARVLAAGSDGEQVSLRLNGHPGCVWANGSAPHERRPNGYALGLVSGGSAQESQGNLGLPDVELLSHELKGVCEPPGLDDPLAQDGAGVS		
rA2a	RIREFRQTFRKIIIRSHVLRQEPFKAAGTSARVLAAGSDGEQVSLRLNGHPGCVWANGSAPHERRPNGYALGLVSGGSAQESQGNLGLPDVELLSHELKGVCEPPGLDDPLAQDGAGVS		
hA2b	RHRDFRYTFHKIISRYLLCQADVKSNGOAGVQFALGVGL		
rA2b	RHRDFRYTFHKIISRYLLCQADVKSNGOAGVQFALGVGL		
rA3	KIKKFKETYFVILRACRLCQTSDSLDSNLEQTTE		
P2Y1	DTFRRRLSRATRKSSRRSEPNVQSKSEMTNLTBYKQNGDTSL		
P2U	QRLVRFARDAKP		

Fig. 2. Deduced amino acid sequences of cloned purinoceptors. The transmembrane helices (TM-1 to TM-7) are designated on the basis of hydrophathy plots. The human A<sub>1</sub> receptor (Salvatore *et al.*, 1992), the rat A<sub>1</sub> receptor (Mahan *et al.*, 1991), the canine A<sub>1</sub> receptor (Libert *et al.*, 1991) and the bovine A<sub>1</sub> receptor (Maenhaut *et al.*, 1990) all have 326 amino acids. The human (Salvatore *et al.*, 1992) and canine (Maenhaut *et al.*, 1990) A<sub>2a</sub> receptors have 412 and the rat A<sub>2a</sub> receptor (Fink *et al.*, 1992) 410 amino acids. The rat (Stehle *et al.*, 1992) and human (Salvatore *et al.*, 1992) A<sub>2b</sub> receptors have 332 amino acids, whereas the rat A<sub>3</sub> receptor (Meyerhof *et al.*, 1991; Zhou *et al.*, 1992) has 319. The P2Y<sub>1</sub> receptor has 362 amino acids (Webb *et al.*, 1993) and the putative P2U receptor has 373 amino acids (Lustig *et al.*, 1993). Modified from Fredholm *et al.* (1994).

*et al.*, 1993) and the 'P<sub>2u</sub>' (Lustig *et al.*, 1993; Parr *et al.*, 1994) purinoceptors is taken into considerations in terms of P2Y purinoceptor subclassification (Section 4.4).

### 3. NEW ATP DERIVATIVES AS POSSIBLE TOOLS TO DISCRIMINATE P2 PURINOCEPTOR SUBTYPES

The confusion surrounding the classification of ATP receptors is partly due to a multiplicity of biological effects and inconsistencies in the potency of available agonists and antagonists (Slinn, 1989; Inoue and Nakazawa, 1992). In this respect, many efforts recently have been aimed at synthesizing novel ATP analogues endowed with high selectivity and potency for the different P2 purinoceptor subtypes.

In collaboration with Burnstock's and Harden's laboratories, Jacobson and coworkers recently have designed ATP derivatives with modified purine, ribose or triphosphate moieties and characterised their activity in a variety of pharmacological assay systems known to possess P2X and P2Y-purinoceptors (Fischer *et al.*, 1993; Burnstock *et al.*, 1994). Pharmacological assays at P2Y purinoceptors included: stimulation of the production of IP3 formation in turkey erythrocytes, relaxation of guinea-pig taenia coli and of rabbit mesenteric artery smooth muscle, and endothelium-dependent relaxation of rabbit aorta. Pharmacological assays at P2X purinoceptors included contraction of the rabbit saphenous artery and contraction of the guinea-pig vas deferens and urinary bladder.

This data is summarised in Table 3, which reports the EC<sub>50</sub> or PD2 values (—log EC<sub>50</sub>) of all the compounds tested in the pharmacological assays. Some analogs displayed selectivity or specificity for either the P2X or P2Y purinoceptor in selective tissues, suggesting heterogeneity within these two purinoceptor families.

Consistent with data showing enhanced potency at P2-purinoceptors of long chain functionalised congeners of 2-mesATP (Zimmet *et al.*, 1993), 2-alkylthio derivatives of ATP proved to be the most potent compounds in all the P2Y-purinoceptor assays (Table 3). Indeed, both base modifications, leading to 8-(6-aminohexylamino)ATP and N-oxide ATP, and ribose modifications, leading to 2,3'-isopropylidene-AMP, resulted in potent derivatives that displayed selectivity for endothelial P2Y purinoceptors, and were practically inactive in the other P2Y- and P2X-purinoceptor pharmacological assays (Table 3). Conversely, the potent agonist N<sup>6</sup>-methyl-ATP and the somewhat less potent agonist 2'-deoxy-ATP were selective for guinea-pig taenia coli P2Y purinoceptors, but were inactive at the vascular P2Y and at the P2X purinoceptors. These results suggest

Table 2. Percentage identity of Chick P2Y<sub>1</sub> Purinoceptor with known G-Protein-Coupled Receptors

Receptor	Species	%Identity with P2Y <sub>1</sub>
RDCI	Canine	27
Angiotensin II type 1	Human	27
Thrombin	Human	25
Platelet activating factor	Guinea-pig	25
CSa anaphylatoxin	Human	23
Neuromedin K	Rat	23
Interleukin 8	Human	22
Bradykinin B2	Rat	22
Neurotensin	Rat	21
Endothelin B	Human	21
Gastrin-releasing peptide	Mouse	21
Adenosine A <sub>1</sub>	Canine	21
Substance P	Human	20
Neurokinin 2	Human	20
Adenosine A <sub>2</sub>	Canine	18
CAMP	Slime mold	17

The most related sequences are shown, along with adenosine and CAMP receptor, RDCI, canine orphan receptor RDCI (Libert *et al.*, 1989). Adapted from Webb *et al.* (1993).



Table 3. Activity of Nucleotide Analogs in Various Biochemical and Pharmacological Models

Compound	Biochemical assay Turkey erythrocyte <sup>1</sup>	P2Y purinoceptors			P2X purinoceptors		
		Mediating relaxation (relative to ATP) <sup>2</sup>			Mediating contraction (relative to ATP) <sup>2</sup>		
		Guinea-pig taenia	Rabbit aorta	Rabbit mesenteric artery	Rabbit saphenous artery <sup>3</sup>	Guinea-pig vas deferens	Guinea-pig bladder
<i>ATP and triphosphate modifications</i>							
1. ATP	2800 ± 700	= (6.2) <sup>4</sup>	= (4.5)	= (6.0)	= [5%]	= (3.5) <sup>5</sup>	= <sup>5</sup>
2. ADP	8000 ± 2000	= <sup>5</sup>	= (5.2)	= (5.2)	na	= <sup>5</sup>	= <sup>5</sup>
3. AMP	4 ± 2% at 10 <sup>-4</sup> M	= <sup>5</sup>	= (4.8)	= (5.0)	na	= <sup>5</sup>	= <sup>5</sup>
4. αβmeATP	>> 100,000	= (5.6) <sup>5</sup>	See footnote 6	See footnote 6	na	= <sup>5</sup>	na <sup>5</sup>
5. βγmeATP	>> 100,000	= <sup>5</sup>	See footnote 6	See footnote 7	= [5.9%]	+ (5.3) <sup>5</sup>	+ + (5.7) <sup>5</sup>
6. βγme-3',5'-cyclic ATP	na	= <sup>5</sup>	na	na	+ + [89%]	+ <sup>5</sup>	+ + <sup>5</sup>
7. AppNHp	4,450 ± 1,150	= <sup>5</sup>	na	na	na	na	na
8. ATPαS (S-isomer)	8,930 ± 4,440	+ <sup>5</sup>	+ (5.5)	+ (6.6)	na	+	+
9. ATPγS	1,260 ± 380	= <sup>5</sup>	See footnote 6	+ (6.6)	na	= <sup>5</sup>	= <sup>5</sup>
10. ADPβS	96 ± 27	=	+ (5.7)	+ (5.8)	na	+ + <sup>5</sup>	+
			+ (5.8)	+ (5.8)	na	= <sup>5</sup>	= <sup>5</sup>
<i>Base modifications</i>							
11a. 2meSATP	8 ± 2	+ + (8.0)	+ + (6.8) <sup>5</sup>	+ + (6.5)	na	= <sup>5</sup>	= <sup>5</sup>
11b. 2-(6-cyano- hexylthio)-ATP	10 ± 5	+ + (8.8)	+ + (6.9)	+ + (7.0)	= [9.2%]	+ +	See footnote 8
12. N <sup>6</sup> methyl-ATP <sup>5</sup>	19,000 ± 6,000	+ (5.8)	=	na	na	na	na <sup>5</sup>
13. 8-Bromo-ATP	47,000	=	na	na	na	na	=
14. 8-(6-Amino- hexylamino)-ATP	8,200 ± 1,200	=	+ + (7.3, < max)	na	na	na	na
15. Adenosine N1-oxide 5'-triphosphate	16,900 ± 4,900	= (4.9)	+ + (6.7, > max)	na	na	na	na
16. N1,N <sup>6</sup> -etheno-ATP	>> 100,000	=	na	na	na	na	na
17. UTP	143,000 ± 44,000	= (3.5)	= (4.8)	+ (6.7)	na	na	= <sup>5</sup>
18. 5-Fluoro-UTP	>> 100,000	=	+ (6.0, ≈ max)	na	na	na	+ +

Table 3—Continued

Compound	Biochemical assay Turkey erythrocyte <sup>1</sup>	P2Y purinoceptors			P2X purinoceptors		
		Mediating relaxation (relative to ATP) <sup>2</sup>			Mediating contraction (relative to ATP) <sup>2</sup>		
		Guinea-pig taenia	Rabbit aorta	Rabbit mesenteric artery	Rabbit saphenous artery <sup>3</sup>	Guinea-pig vas deferens	Guinea-pig bladder
<i>Ribose modifications</i>							
19. 2'-Deoxy-ATP	19,200 ± 6,200	= 5.6	na	na	na	na <sup>5</sup>	na
20. 3'-Deoxy-ATP	75,500 ± 14,800	—	na	na	na	=	—
21. 2',3'-Dideoxy-ATP	70,800	= (5.0)				=	+
22. 3'-Amino-3'- deoxy-ATP <sup>5</sup>	193	= (5.4)	++ (6.4, ≈ max)	na	na	+	=
23. 3'-Acetylamino- 3'-deoxy-ATP <sup>5</sup>	» 100,000	na	na	+ (< max)	na	=	+
24. 3'-(4-Hydroxyphenylpro pionylamino)-3'-deoxy-ATP <sup>5</sup>	» 100,000	na	++ (< max)	na	na	=	+
25. 3'-Benzylamino-3'- deoxy-ATP	» 100,000	—	na	na	na	++	++
26. Isopropylidene-ATP	201,000 ± 63,000	—	na	—	= [7.7%]	na	—
27. Isopropylidene-AMP <sup>5</sup>	» 100,000	na	++ (< max)	na	na	na	na

<sup>1</sup> EC<sub>50</sub> values (nM) for stimulation of production of inositol phosphates, expressed as the mean ± S.D. for at least 3–5 determinations, or % stimulation at concentration indicated; other symbols as in footnote 2. <sup>2</sup> + +, significantly more potent than ATP; +, more potent than or equal to ATP; =, equal to ATP; —, less potent than or equal to ATP; — —, significantly less potent than ATP; na, not active at the highest concentration tested (usually around 10<sup>-3</sup> M). Numerical value, if given, is pD<sub>2</sub> in minus log units, and for some compounds in rat aorta and mesenteric artery, maximum relaxation relative to 2meSATP (<,  $\approx$ , or >) is indicated in parentheses. <sup>3</sup> For the saphenous artery, the percentages in brackets are responses at 10  $\mu$ M relative to the contraction produced by 1  $\mu$ M  $\alpha$ meATP. The highest concentrations tested were 3–10  $\mu$ M. <sup>4</sup> 6.2 ± 0.08 (n = 38). <sup>5</sup> Data for smooth muscle from literature reports (Burnstock *et al.*, 1983, 1984; Cooper *et al.*, 1989; Cusack and Hourani, 1990; Jacobson, 1990). <sup>6</sup> Contraction, not relaxation. <sup>7</sup> Relaxation, but pD<sub>2</sub> not calculable. <sup>8</sup> Compound 11b was approximately 100 times more potent than ATP in the bladder, but it produced tonic contractions rather than the phasic contractions of ATP. In the presence of indomethacin (1  $\mu$ M), it was much less potent than ATP under the same conditions. Modified from Burnstock *et al.* (1994).



P2Y-purinoreceptor heterogeneity, with purinoreceptors in the taenia coli and those in the vasculature being recognised differently by ATP derivatives carrying selective structural modifications.

Similarly, heterogeneity of P2X purinoreceptors is also suggested. Ribose modifications leading to 3-benzylamino-3'-deoxyATP resulted in a very potent agonist at the P2X purinoreceptors in guinea-pig vas deferens and bladder, with no activity in rabbit saphenous artery and at P2Y purinoreceptors; this would suggest that the vascular P2X purinoreceptor differs from vas deferens and bladder P2X purinoreceptors, which appeared to be very similar to each other. This conclusion is also supported by the fact that a number of synthesised derivatives displayed no activity at the saphenous artery P2X purinoreceptor, but could not distinguish between the vas deferens and urinary bladder P2X purinoreceptors (Table 3). Interestingly, the only compound distinguishing between the vas deferens and the bladder P2X purinoreceptor was the pyrimidine compound 5-fluoro-UTP. Whether this is indicative of a true pyrimidine receptor or whether the P2X purinoreceptor in the bladder also recognises pyrimidine compounds has yet to be evaluated.

It is apparent from the structure-activity data given in Table 3 that the P2 purinoreceptors in the seven tested tissues are all different from one another in their pharmacological profile. This would suggest, therefore, that within the P2X- and P2Y-purinoreceptor families there are further subtypes that can be distinguished by the selective actions of ATP derivatives.

#### 4. NEW PROPOSALS FOR P2-PURINORECEPTOR CLASSIFICATION

##### 4.1. P2-Purinoreceptor Coupling to Different Transduction Mechanisms: A Criterion for Distinguishing the P2X-(Ligand-Gated) and P2Y-(G-Protein-Coupled) Purinoreceptor Families

There is now good evidence to suggest that the P2X- and P2Y-purinoreceptor subtypes differ not only in terms of pharmacological profile and tissue distribution, but also in their transduction mechanisms and effector systems. We propose, therefore, the existence of P2X- and P2Y-purinoreceptor families, each with their own subdivisions.

Evidence to date suggests that P2X purinoreceptors are intrinsic ionic channels permeable to Na<sup>+</sup>, K<sup>+</sup> and Ca<sup>2+</sup>, whereas P2Y purinoreceptors are G-protein-coupled receptors. A common feature of the P2Y-purinoreceptor family is the activation, following receptor binding by selective agonists, of a membrane shuttle protein belonging to the GTP-binding protein family (G-proteins), which, in turn, activates selective enzyme effector systems that regulate intracellular concentrations of second messengers (for a review, see Gilman, 1987). In the case of P2Y purinoreceptors, effector systems are represented by phospholipase C (which modulates IP<sub>3</sub> and DAG formation), phospholipase A<sub>2</sub> (with consequent generation of arachidonic acid metabolites), or adenylate cyclase (which modulates cAMP levels) (Fredholm *et al.*, 1994). There is now sufficient data to suggest a further subdivision of P2X and P2Y purinoreceptors into subclasses (see Section 3), and based on this, we can hypothesise the existence of P2X-(ligand-gated) and P2Y-(G-protein-coupled) purinoreceptor families.

##### 4.2. Do the P<sub>2u</sub>, P<sub>2o</sub> and P<sub>2s</sub> Receptors Belong to the P2Y-Purinoreceptor Family?

As reported in Table 1, the so-called P<sub>2u</sub>, P<sub>2o</sub> and the P<sub>2u</sub>/P<sub>2o</sub>'pyrimidine' receptors share with 'classic' P2Y purinoreceptors the characteristic of utilizing G-proteins for their transduction mechanisms.

Based on this and other evidence, we would like to propose the inclusion of these purinoreceptor subtypes within the P2Y-purinoreceptor family.

All present and yet-to-be identified receptors belonging to this P2Y-purinoreceptor family would involve G-protein activation, and would be named P2Y<sub>1</sub>, P2Y<sub>2</sub>, P2Y<sub>3</sub>, ... P2Y<sub>n</sub>, with the clear advantages of eliminating the confusion generated by the 'D', 'V' and 'U' subheadings and of easily accommodating any 'new' P2Y-purinoreceptor subtype by simply assigning it a progressive number. Apart from the G-protein involvement, there are a number of additional reasons to support this new way of defining the different ATP receptor subtypes.

In the case of the P<sub>2u</sub>/P<sub>2o</sub>'nucleotide' receptor, such a classification would be supported by the fact that this receptor subtype recognises both UTP and ATP and has in common with the recently

#### Purinoreceptors: Are there families of P2X and P2Y purinoreceptors?

Table 4. Proposed Subclassification of P2 Purinoreceptors

Name	P2X-purinoreceptor family	P2Y-purinoreceptor family <sup>1</sup>	P2Z
Type	Ligand-gated channel	G-protein-coupled	Non selective pore
General agonist profile	$\alpha\beta\text{meATP} > \beta\text{meATP} > \text{ATP} \approx \text{2meSATP} \approx \text{ADP}$	$\text{2meSATP} > \text{ATP} = \text{ADP}$	ATP <sup>+</sup>
Antagonists	$\alpha\beta\text{meATP}$ desensitisation Suramin Selectively blocked by PPADS ANAPP3	Suramin Reactive blue 2	Oxidised ATP

<sup>1</sup>It is proposed that the P2Y family also includes the former G-protein-linked P<sub>2u</sub>/P<sub>2o</sub>'pyrimidine' receptor (where UTP  $\geq$  ATP), the P<sub>2s</sub> purinoreceptor (where ADP is the unique agonist) and the P<sub>2o</sub>-purinoreceptor (where APx<sub>A</sub> represents the specific agonists: see text for details).

cloned P2Y<sub>1</sub>-purinoreceptor sequences encoding for the same group of G-protein-coupled receptors (Section 2).

For the P<sub>2s</sub> receptor, inclusion into the P2Y-purinoreceptor family as P2Y<sub>3</sub> is proposed following the isolation of a related receptor (clone 103, Barnard *et al.*, 1994) during the cloning work with chick brain, which led to the isolation of the P2Y<sub>1</sub> purinoreceptor (clone 803, Webb *et al.*, 1993).

For the P<sub>2o</sub> receptor subtype, inclusion into the P2Y-purinoreceptor family would be justified by the fact that, although the major agonists of this subclass are dinucleotides rather than mononucleotides, this receptor subtype displays a 'P2Y-like' pharmacological profile in chromaffin cells and *Torpedo* synaptic terminals (Pintor and Miras-Portugal, 1993). As summarised in Table 4, acquisition of this new nomenclature would lead to a simpler purinoreceptor classification comprising the major divisions P2X/P2Y/P2Z.

To support our hypothesis, we have analyzed carefully a number of papers published between 1987 and 1994 on the pharmacological actions of ATP and its analogues. Particular attention has been paid to comparative studies where, besides ATP itself, a variety of different ATP analogues have been tested, in an attempt to assess whether specific pharmacological profiles could be defined in different tissues and systems.

##### 4.3. Proposals for the Subclassification of the P2Y-Purinoreceptor Family

By carefully analysing the studies published so far on P<sub>2</sub>-like responses, we were able to define at least seven different agonist potency profiles, which, indeed, would suggest the existence of seven P2Y-purinoreceptor subclasses (Table 5).

Two major considerations need to be taken into account before describing in detail the proposed P2Y-purinoreceptor subtypes. First, examples listed in Table 5 only refer to cases in the literature where all the key agonists selected by us to establish the rank order of potency for the putative P2Y<sub>1-7</sub> purinoreceptor subtypes have been tested. Of course, there are many more examples of P2Y<sub>1-7</sub> responses in a variety of other tissues or systems not listed in Table 5, where, however, only some of the key compounds have been tested, therefore making it impossible to 'assign' the tissue to a specific receptor subtype. Second, the reader might notice that some tissues are listed twice under two different P2Y-purinoreceptor subtypes. Whether this is due to the co-presence of multiple purinoreceptor populations in the same tissues or to different tissue preparations and/or animal species is difficult to say at the moment. The use, where available, of molecular probes for the P2Y-purinoreceptor subtypes and of the new selective P2Y agonists (Section 3) will be of help in clarifying this aspect.

The P2Y<sub>1</sub> receptor recently cloned by Webb *et al.* (1993) shows the following agonist potency profile: 2meSATP  $\geq$  ATP  $\gg$  ADP  $\gg$   $\alpha\beta\text{meATP}$ . We were able to find several examples of systems displaying this agonist potency profile (see list in Table 5). In some cases, pharmacological data correlated well with expression of the chick P2Y<sub>1</sub> transcript, as determined in Webb and coworkers' study by Northern hybridisation. This would suggest strongly that the clone identified by Webb *et al.* is the P2 purinoreceptor independently studied by other researchers showing this typical agonist

Table 5. Proposed Subclassification of the P<sub>2</sub>Y G-Protein-Linked Purinoceptor Family

Proposed subtype	P <sub>2</sub> Y <sub>1</sub>	P <sub>2</sub> Y <sub>2</sub>	P <sub>2</sub> Y <sub>3</sub> †	P <sub>2</sub> Y <sub>4</sub>	P <sub>2</sub> Y <sub>5</sub>	P <sub>2</sub> Y <sub>6</sub>	P <sub>2</sub> Y <sub>7</sub> ‡
Agonist potency	2meSATP ≥ ATP > ADP > αβmeATP	ATP ≥ UTP = ATPγS; UTP > ATP only for footnotes 1-3	2meSADP > ADP	2meSATP > ATP = ADP = αβmeATP > βγmeATP	2meSATP ≥ ATP = ADP > αβmeATP βγmeATP inactive	2meSATP > ATP > ADP?	Diadenosine polyphosphates
Selective agonists	2meSATP	UTPγS		2'-deoxy-ATP and also N <sup>6</sup> -methyl-ATP selective for <i>taenia coli</i>	8-(6-aminohexylamino)-ATP and ATP-N-oxide selective for <i>endothelial cells</i>	No selective agonists available yet; however, the compounds selective for P <sub>2</sub> Y <sub>4</sub> and P <sub>2</sub> Y <sub>5</sub> are inactive on <i>vascular smooth muscle</i>	
Name of clone	Clone 803 (Webb <i>et al.</i> , 1993)	Clone pP2R (Lustig <i>et al.</i> , 1993) Clone HP2U (Parr <i>et al.</i> , 1994) Clone from rat heart tissue, (Godecke and Schrader, 1994)	Brain-derived clone 103 <sup>4</sup>				
Number of amino acids	362	373 for clone pP2R 375 for clone HP2U					

Proposed subtype	P <sub>2</sub> Y <sub>1</sub>	P <sub>2</sub> Y <sub>2</sub>	P <sub>2</sub> Y <sub>3</sub> †	P <sub>2</sub> Y <sub>4</sub>	P <sub>2</sub> Y <sub>5</sub>	P <sub>2</sub> Y <sub>6</sub>	P <sub>2</sub> Y <sub>7</sub> ‡
Tissue	Chick brain <sup>5</sup> Rat brain cortex <sup>6</sup> Developing chick skeletal muscle <sup>5,7</sup> Guinea-pig cochlear hair cells <sup>8</sup> Rabbit gastric glands <sup>9</sup>	Rat renal mesangial cells <sup>10,11</sup> Rat mesenteric arterial bed <sup>12</sup> Rat aortic smooth muscle cells <sup>13</sup> Rat liver <sup>11,14</sup> Rat hepatocytes <sup>11,15</sup> Rat osteoblastic cells <sup>16</sup> Rabbit <sup>17</sup> and human <sup>18</sup> neutrophils Guinea-pig brain <sup>19</sup> Sheep pituitary cells <sup>20</sup> Rat myocardial endothelium <sup>1,11</sup> and bovine endothelial cells <sup>21</sup> Skeletal muscle <sup>11</sup> Human fibroblasts <sup>22</sup> Human amnion cells <sup>2,11</sup> Human airway epithelial cells <sup>11,23</sup> HL60 cells <sup>24-26</sup> PC12 cells <sup>27</sup> Mouse neuroblastoma cells <sup>28</sup>	Platelets <sup>29</sup> Megakaryocytes <sup>30</sup> Brain capillary endothelial cells <sup>31</sup>	Guinea-pig taenia coli <sup>3</sup> Turkey erythrocytes <sup>32-33</sup>	Rabbit aorta <sup>35</sup> Pig aorta <sup>36</sup> Rat cortical astrocytes <sup>37,38</sup> Organ of Corti cells <sup>39</sup> Dorsal spinal cord astrocytes <sup>40</sup>	Rabbit coronary <sup>41</sup> , hepatic <sup>42</sup> and mesenteric <sup>43</sup> arteries Human subcutaneous and omental resistance arteries <sup>44</sup> Human endothelial cells <sup>44</sup>	Rat brain synaptosomes <sup>45</sup> Rat cortical neurons <sup>46</sup> Chromaffin cells <sup>47-49</sup> Porcine <sup>50</sup> and bovine <sup>51</sup> aortic endothelial cells Human platelets <sup>52-54</sup> Mammalian hepatocytes <sup>55</sup> Guinea-pig vas deferens, urinary bladder <sup>56</sup> and isolated arteries <sup>57</sup>

<sup>1</sup>Formerly P<sub>2</sub> or P<sub>2</sub> or 'nucleotide' receptor. <sup>†</sup>Formerly P<sub>2</sub>. <sup>‡</sup>Formerly P<sub>2</sub>D. <sup>4</sup>As suggested by hybridisation studies with rat heart mRNA (Godecke and Schrader, 1994). <sup>5</sup>Vander Kooy *et al.* (1989). <sup>6</sup>Satchell and Maguire (1975). <sup>7</sup>Maguire and Satchell (1979). <sup>8</sup>Burnstock *et al.* (1983). <sup>9</sup>Cusack *et al.* (1987). <sup>10</sup>Two quite different full-length cDNAs were isolated by Webb *et al.* (1993); one encoding for the P<sub>2</sub>Y<sub>1</sub> purinoceptor, the other encoding for a receptor responsive to ADP (Barnard *et al.*, 1994). Preliminary data suggest that this second recombinant brain-derived receptor is similar, but not identical, to the P<sub>2</sub> receptor of mammalian platelets. This is also consistent with Frelin *et al.* (1993). <sup>11</sup>Also supported by hybridisation with P<sub>2</sub>Y<sub>1</sub> mRNA (Webb *et al.*, 1993). <sup>12</sup>von Kugelgen *et al.* (1994); first description of presynaptic P<sub>2</sub> purinoceptors in the CNS. <sup>13</sup>Thomas *et al.* (1991). <sup>14</sup>Nakagawa *et al.* (1990). <sup>15</sup>Gil-Rodrigo *et al.* (1990). <sup>16</sup>Reactive blue 2-dependent effect; clear evidence for G-protein involvement; no evidence for separate ATP and UTP receptors; Pfeilschifter (1990). <sup>17</sup>As shown by hybridisation with HP2U mRNA (Parr *et al.*, 1994). <sup>18</sup>Ralevic and Burnstock (1991). <sup>19</sup>Pertussis-toxin sensitive effect (Tawada *et al.*, 1987). <sup>20</sup>Haussinger *et al.* (1987). <sup>21</sup>Keppens *et al.* (1992). <sup>22</sup>Clear evidence for a common UTP/ATP receptor-mediated release of Ca<sup>2+</sup> from internal stores; these cells also have a receptor to ADP (Reimer and Dixon, 1992). <sup>23</sup>Elferink *et al.* (1992). <sup>24</sup>Pertussis toxin-sensitive effect (Walker *et al.*, 1991). <sup>25</sup>As shown in *Xenopus* oocytes injected with guinea-pig brain mRNA (Honore *et al.*, 1991). <sup>26</sup>Phospholipase C-mediated effects (Davidson *et al.*, 1990). <sup>27</sup>Phospholipase C-mediated effects; common UTP/ATP receptor suggested (Purkiss *et al.*, 1993). <sup>28</sup>Fine *et al.* (1989). <sup>29</sup>Evidence for a common ATP/UTP receptor coupled to phospholipase C (Mason *et al.*, 1991; Brown *et al.*, 1991); nature of this receptor has been confirmed recently by functional expression of the cloned human receptor in 1321N1 cells (Lazarowski *et al.*, 1994). <sup>30</sup>Xing *et al.* (1992). <sup>31</sup>Cockcroft and Stutchfield (1989). <sup>32</sup>Also shown in *Xenopus* oocytes injected with mRNA from HL60 cells (Murphy and Tiffany, 1990). <sup>33</sup>Evidence for an ATP/UTP phospholipase-linked receptor (Murrin and Boarder, 1992); however, it is not clear whether multiple receptors for ATP and UTP are expressed in these cells (Reed *et al.*, 1994). <sup>34</sup>As shown by hybridisation with pP2R mRNA (Lustig *et al.*, 1993). <sup>35</sup>Cusack *et al.* (1979). <sup>36</sup>Burnstock (1991). <sup>37</sup>ADP-specific receptor inducing Ca<sup>2+</sup> mobilisation from a thapsigargin-sensitive intracellular pool (Frelin *et al.*, 1993). <sup>38</sup>Berrie *et al.* (1989). <sup>39</sup>Boyer *et al.* (1990). <sup>40</sup>2-(6-cyanoheptylthio)-ATP shows nanomolar potency at the turkey erythrocyte P<sub>2</sub>Y receptor (Fischer *et al.*, 1993). <sup>41</sup>Burnstock *et al.* (1994). <sup>42</sup>Martin *et al.* (1985). <sup>43</sup>Phospholipase C-linked receptor (Kastritis *et al.*, 1992). <sup>44</sup>Activation of this receptor leads to intracellular Ca<sup>2+</sup>-dependent arachidonic acid release (Bruner and Murphy, 1990). <sup>45</sup>Dulon *et al.* (1993). <sup>46</sup>Activation of this receptor leads to mobilisation of Ca<sup>2+</sup> from a thapsigargin-sensitive intracellular store (Salter and Hicks, 1994). <sup>47</sup>Brizzolara and Burnstock (1991). <sup>48</sup>αβmeATP practically inactive (Martin *et al.*, 1991). <sup>49</sup>Carter *et al.* (1988) reported that activation of this receptor on human cultured umbilical vein endothelial cells induces intracellular Ca<sup>2+</sup>-dependent prostacyclin secretion. Evidence was produced in favour of the different nature of this receptor from the taenia coli P<sub>2</sub>Y purinoceptor. <sup>50</sup>Pintor *et al.* (1993). <sup>51</sup>Stone and Perkins (1981). <sup>52</sup>Pintor *et al.* (1991a). <sup>53</sup>Pintor *et al.* (1991b). <sup>54</sup>Castro *et al.* (1992). <sup>55</sup>Goldman *et al.* (1986). <sup>56</sup>Ogilvie *et al.* (1989). <sup>57</sup>Flodgaard and Klenow (1992). <sup>58</sup>Luthje and Ogilvie (1983). <sup>59</sup>Zamecnik *et al.* (1992). <sup>60</sup>Rapaport and Zamecnik (1976). <sup>61</sup>Hoyle *et al.* (1989); Hoyle (1990). <sup>62</sup>Busse *et al.* (1988).

potency profile. In other cases, there was a clear, although not yet conclusive, matching between the pharmacological data and the results of the hybridization studies with P2Y<sub>1</sub> mRNA. For example, Thomas *et al.* (1991) reported the presence of a P2 purinoceptor (whose pharmacological profile was not unequivocally assessed) in chick skeletal muscle, a tissue that was shown to abundantly express the P2Y<sub>1</sub> transcript. However, it is not clear if the receptor described by Thomas *et al.* is a G-protein-linked receptor or a ligand-gated cation channel.

In analysing the abundant literature on the UTP-sensitive receptor, we could identify two receptor behaviour patterns: in most cases, UTP was equipotent with ATP, whereas in a few other examples, UTP was more potent than ATP. At present, there is not sufficient data to support the existence of two different UTP-sensitive purinoceptors (the first one preferentially responding to ATP and the second one preferentially responding to UTP), and we, therefore, put together the two agonist behaviour patterns and named this purinoceptor P2Y<sub>2</sub> (Table 5). UTP<sub>2</sub>S seems to behave as a selective agonist at this receptor subtype (Table 5).

Based on our proposal, clone pP2R (Lustig *et al.*, 1993), therefore, would be the P2Y<sub>2</sub>-purinoceptor subtype. The detection of the pP2R transcript in kidney, liver, lung, heart and brain (Lustig *et al.*, 1993) matches well with the reported presence of the P2Y<sub>2</sub> purinoceptor in rat renal mesangial cells (Pfeilschifter, 1990), rat hepatocytes (Keppens *et al.*, 1992), human airway epithelial cells (Mason *et al.*, 1991; Brown *et al.*, 1991; Lazarowski *et al.*, 1994) and guinea-pig brain (Honore *et al.*, 1991; Table 5). As underlined in Table 5, a good correlation from previous literature reports and expression of receptor mRNA was also found for the recently cloned HP2U purinoceptor (Parr *et al.*, 1994). In the examples listed in Table 5, the effects of ATP and UTP do not seem to be additive when both agonists are utilised at maximal doses suggesting the presence of a common ATP/UTP receptor (references 10, 15, 16 and 23 of Table 5), in agreement with the conclusion recently drawn by Keppens (1993) in reviewing the effects of ATP and UTP in isolated hepatocytes. Moreover, where tested, UTP responses appear to be antagonised by reactive blue 2 or suramin, which are known to act as antagonists at 'classic' P<sub>2</sub> purinoceptors (Hoyle *et al.*, 1990). Both these characteristics further strengthen the inclusion of this receptor into the P<sub>2</sub>Y-purinoceptor family.

Nonetheless, there may be a need to recognise new classes of receptor associated with the P2 families that are not yet included in our proposed classification. For example, recent data from the C6-2B rat glioma cell line identified a phospholipase C-linked receptor, which was claimed to be selectively activated by UTP and related uridine nucleotides, but showed insensitivity to ATP, thus differentiating it from the pP2R and HP2R clones (Lazarowski and Harden, 1994). This receptor may represent a new class of 'P<sub>2</sub>-like' receptor. For the long term, the resolution of whether there are distinct P2Y<sub>2</sub> ('P2U') and 'UTP' (pyrimidine) receptors will be dependent on the generation of new data, both molecular and pharmacological, that will unambiguously delineate the nature of the receptor subtypes responding to UTP (Williams, 1994).

The currently named P<sub>2</sub> receptor (P2Y<sub>3</sub> in our proposed subclassification) seems to be expressed exclusively by platelets and megakaryocytes. However, we were able to find a very recent report of a similar ADP receptor in brain capillary endothelial cells (Frelin *et al.*, 1993), which would suggest a wider biological role for this receptor subtype. The possible presence in brain of a P2-purinoceptor subtype displaying a strong preference to ADP is also suggested by the second recombinant brain-derived receptor recently cloned by the Barnard/Burnstock London group and obtained during the polymerase chain reaction amplification of sequences encoding G-protein-coupled receptors, which led to the cloning of the P2Y<sub>1</sub>-purinoceptor subtype (Webb *et al.*, 1993). This second clone, designated P2Y<sub>1</sub>, is largely similar (although not identical) to the P<sub>2</sub> purinoceptor of mammalian platelets, and Northern hybridisation studies have revealed the presence of this mRNA in brain and several peripheral tissues (Barnard *et al.*, 1994). These results confirm that the currently named P<sub>2</sub> receptor also belongs to the G-protein-coupled receptor superfamily and underlines its close relationship to the P2Y-purinoceptor family.

In our analysis of the literature in this area, we also found examples in favour of three additional agonist potency profiles: 2mSsATP ≫ ATP = ADP ≫ αβmeATP ≫ βγmeATP (P2Y<sub>4</sub>), 2mSsATP ≫ ATP = ADP ≫ αβmeATP (βγmeATP inactive) (P2Y<sub>5</sub>), and 2mSsATP > ATP > ADP (P2Y<sub>6</sub>). These three additional receptors can now be discriminated from each other with the aid of newly synthesised compounds.

The prototype for the P2Y<sub>4</sub> subtype is represented by the guinea-pig taenia coli, and this receptor

seems to be selectively activated by the new compounds 2'-deoxy-ATP and N<sup>6</sup>-methyl-ATP (Table 5 and Section 3). A characteristic of this receptor is the much higher sensitivity to 2mSsATP with respect to ATP. For example, 2mSsATP is 250-fold more potent than either ATP or ADP in stimulating phosphatidylinositol breakdown in turkey erythrocytes (Bertie *et al.*, 1989).

The prototype for the P2Y<sub>5</sub> subtype is represented by the endothelial receptor mediating blood vessel relaxation via generation of nitric oxide. The new compounds 8-(6-aminohexylamino)-ATP and ATP-N-oxide, which were shown to be selective for aortic endothelial cells (Burnstock *et al.*, 1994), could behave as selective P2Y<sub>5</sub> purinoceptor agonists and will be useful tools to be tested in the biological systems displaying a similar agonist potency profile (Refs 37-40 of Table 5), which, besides endothelial cells, also include astrocytes and Organ of Corti cells.

The P2Y<sub>6</sub> purinoceptor is the vascular smooth muscle receptor responsible for direct ATP vasodilation (Refs 35, 41 and 42 of Table 5). No selective agonists are available yet for this purinoceptor subtype. However, derivatives selective for the P2Y<sub>4</sub> and P2Y<sub>5</sub> subtypes are inactive at this receptor subtype. Again, future studies aimed at testing the activity of the new compounds in a variety of tissues will confirm the identity and nature of this hypothetical purinoceptor subtype. Interestingly, one of the newly synthesised analogs, 2-(6-cyanoheptylthio)-ATP, was proven to be extremely potent at both the guinea-pig taenia coli and the rabbit aorta and the rabbit mesenteric artery (P2Y<sub>4</sub>-P2Y<sub>6</sub>) purinoceptor subtypes (Fischer *et al.*, 1993; Burnstock *et al.*, 1994). This compound belongs to the family of long-chain 2-alkylthio-ATP derivatives, which were previously shown to resist degradation by nucleotidases (Zimmet *et al.*, 1993) and may serve as the basis for the design of useful molecular probes for ATP receptors.

Finally, an additional purinoceptor subtype included in this receptor family is the P2Y<sub>7</sub> purinoceptor (formerly P<sub>2D</sub>), the presence of which has been demonstrated in a variety of systems (Refs 45-57 of Table 5). The reasons for assigning this receptor to the P2Y<sub>7</sub>-purinoceptor family have been explained previously (Section 4.2) and, again, a definite confirmation, or not, that this represents a P2Y<sub>7</sub>-purinoceptor subtype distinct from the others will come from cloning data.

#### 4.4. Proposals for the Subclassification of the P2X-Purinoceptor Family

The use of new ATP derivatives with modified purine, ribose or triphosphate moieties suggests heterogeneity within the P2X-ion-gated purinoceptors (Burnstock *et al.*, 1994). This heterogeneity is also consistent with the results of our analysis of a series of literature papers reporting P2X-like responses. Again, the references reported here refer to studies where a number of key agonists have been tested and, therefore, this review cannot be comprehensive for all the papers published on P2X-purinoceptor-mediated effects.

Although evidence for subclassification in this case was less obvious than that of the P2Y purinoceptor, there are indications for at least four different agonist potency profiles, leading to the identification of four P2X-purinoceptor subclasses (Table 6).

Guinea-pig vas deferens seems to be the prototype of the first P2X purinoceptor subtype. In general, at this receptor, αβmeATP is equipotent with ATP in inducing contraction (Burnstock *et al.*, 1994; Wiklund and Gustafsson, 1988). This receptor recently has been reported to have been cloned by the Glaxo group (A. Surprenant, personal communication\*), and, therefore, it has been named P2X<sub>1</sub>. It will be, of course, extremely interesting to test the new ATP derivatives shown to be selective for vas deferens (Section 3) on the cloned transfected receptor. Useful information is likely to come in the near future from an analysis of the distribution of its mRNA in the other tissues showing P2X-like responses.

The pharmacology of the urinary bladder P2X purinoceptor appears to be different from that of the vas deferens receptor. In both guinea-pig (Burnstock *et al.*, 1994; Cusack, 1993) and rat urinary bladder (Bo *et al.*, 1994) both αβmeATP and βγmeATP are significantly more potent than ATP and 2mSsATP. The different nature of this receptor is also confirmed by the fact that it can be discriminated from the vas deferens receptor with the new derivative 5-fluoro-UTP (Section 3; Table 6). Both this compound and the other new derivatives reported in Tables 3 and 6 will be invaluable experimental tools to test the presence of this receptor subtype (named P2X<sub>2</sub>) in a number of systems.

\*See Valera *et al.* (1994) in "Note added in proof".



Table 6. Proposed Subclassification of the P2X-Ion Gated Purinoceptor Family

Proposed subtype	P2X <sub>1</sub>	P2X <sub>2</sub>	P2X <sub>3</sub>	P2X <sub>4</sub>
Selective agonists	2-(4-nitrophenylethyl-thio)ATP > 3'-amino-3'-deoxy-ATP = 2-hexyl-thioATP = 2-cyclo-hexyl-thioATP	5-fluoro-UTP > 2-hexyl-thioATP = 3'-acetyl-amino-3'-deoxy-ATP > 3'-(4-hydroxy-phenylpropionyl-amino)-3'-deoxy-ATP	No selective agonists available; however, the compounds selective on the other subtypes are inactive	No selective agonists available
Tissues	Guinea-pig vas deferens <sup>1-3</sup>	Guinea-pig <sup>14</sup> and rat <sup>3</sup> urinary bladder Rabbit bladder detrusor <sup>8</sup> Rat colon longitudinal muscle <sup>7</sup>	Vascular smooth muscle <sup>1</sup>	Peripheral <sup>8-12</sup> and central <sup>13-16</sup> neurons Ventricular myocytes <sup>17</sup> Microglial cells <sup>18</sup>

<sup>1</sup>Burnstock *et al.* (1994), <sup>2</sup>Wiklund and Gustafsson (1988), <sup>3</sup>A. Surprenant, personal communication, <sup>4</sup>Cusack (1993), <sup>5</sup>Bo *et al.* (1994), <sup>6</sup>Kishii *et al.* (1992), <sup>7</sup>Bailey and Hourani (1992), <sup>8</sup>Bean (1990), <sup>9</sup>Allen and Burnstock (1990), <sup>10</sup>Krishtal *et al.* (1983), <sup>11</sup>Fieber and Adams (1991), <sup>12</sup>Surprenant (1994), <sup>13</sup>Ueno *et al.* (1992), <sup>14</sup>Edwards *et al.* (1992), <sup>15</sup>Tschöpl *et al.* (1992), <sup>16</sup>Shen and North (1993), <sup>17</sup>Björnsson *et al.* (1989), <sup>18</sup>Langosch *et al.* (1994).

A third P2X-purinoceptor subclass (P2X<sub>3</sub>) might be represented by the receptor expressed in vascular smooth muscle. At the saphenous artery P2X purinoceptor, 2mS-ATP and the new derivatives selective for the P2X<sub>1</sub> and P2X<sub>2</sub> are inactive (Burnstock *et al.*, 1994; Table 6). It would be interesting to assess the similarities of this receptor with that described by O'Connor *et al.* (1990) in rabbit ear artery. No selective agonists for this receptor subtype are available yet.

An additional P2X-purinoceptor subtype (P2X<sub>4</sub>) might be represented by the neuronal ATP-activated cation channel described in peripheral and central neurons (references 8-16 in Table 6). The general order of potency for ATP analogues in activating this receptor is 2mS-ATP > ATP >  $\alpha$ beta-ATP (Illes and Norenberg, 1993). A similar behaviour is displayed by ATP derivatives also in ventricular myocytes (reference 17 in Table 6) and rat microglial cells (reference 18 in Table 6). The different nature of this receptor from the other hypothesised P2X-purinoceptor subtypes is confirmed by preliminary results, suggesting that the transfected P2X<sub>4</sub> purinoceptor clone behaves differently in terms of pharmacological response from the neuronal ATP-sensitive channel (A. Surprenant, personal communication).

Finally, there are a number of reports in favour of the existence of excitatory purinoceptors responding to both ATP and UTP (von Kugelgen *et al.*, 1987, 1990; von Kugelgen and Starke, 1990; Saig *et al.*, 1990, 1992; Theobald, 1992; Hourani *et al.*, 1993; Pavenstadt *et al.*, 1991; Rubino and Burnstock, 1994; Ralevic and Burnstock, 1991). This receptor differs from the proposed P2Y<sub>2</sub> purinoceptor (= P<sub>2u</sub> purinoceptor), since apparently it is not coupled to G-proteins, but mediates rapid and transient responses (usually excitatory). In some instances and tissues, evidence was produced in favour of a common ATP/UTP receptor (von Kugelgen *et al.*, 1990; Hourani *et al.*, 1993; Ralevic and Burnstock, 1991); in other cases, distinct UTP receptors were hypothesised (Saig *et al.*, 1990, 1992; von Kugelgen *et al.*, 1987). Whether this receptor really represents a distinct entity with respect to the other P2X-purinoceptor subtypes cannot be specified at this time, also because UTP was not tested in all the other studies reported for receptors P2X<sub>1</sub>-P2X<sub>4</sub>. Here again help will come from molecular cloning studies.

We are aware of the fact that differences of agonist order potency within different tissues may partially derive from different ecto-ATPase activities (Cusack, 1993). In this respect, useful and crucial information in terms of purinoceptor classification could come either from the use of dephosphorylation-resistant agonists (e.g. L- $\beta$ -methylene-ATP; Cusack, 1993) or by utilising inhibitors of nucleotide degradation (Leff *et al.*, 1994; Ziganshin *et al.*, 1994a).

## 5. P2-PURINOCEPTOR LIGANDS AS POTENTIAL THERAPEUTIC ENTITIES

While there have been several excellent reviews on the potential development of P1-purinoceptor targeted drugs (Daly, 1982; Daval *et al.*, 1991; Jacobson *et al.*, 1992; Stone, 1992; Williams, 1993), there has been very little discussion on the possible clinical and therapeutic applications of P2-purinoceptor agonists and antagonists (Burnstock, 1993).

Due to the recent explosion of interest in P2 purinoceptors, which has enormously advanced our knowledge on the functional roles of ATP in many different organs and systems, we now have enough experimental data to suggest that P2 purinoceptors might also represent a novel target for drug development in a variety of pathological conditions.

Moreover, the large heterogeneity of P2-purinoceptor subtypes in the different tissues underlined in this review opens the possibility of developing highly selective organ- or tissue-specific P2-purinoceptor-targeted drugs. Several exciting possibilities are being explored.

### 5.1. Diabetes

A number of studies has demonstrated the presence of P2Y purinoceptors on insulin-secreting pancreatic  $\beta$ -cells (Bertrand *et al.*, 1987; Arkhammar *et al.*, 1990; Li *et al.*, 1991; Hillaire-Buys *et al.*, 1992, 1993). The activation of this receptor leads to potent phospholipase C-mediated intracellular Ca<sup>2+</sup> mobilisation, sustained insulin secretion and notable improvement of glucose tolerance. Interestingly, not only was the ecdonucleotide-resistant P2Y-purinoceptor agonist ADP $\beta$ S demonstrated to be effective after oral administration (Hillaire-Buys *et al.*, 1993), but this agent also retained its insulin-stimulatory effects in streptozotocin-diabetic rats (Hillaire-Buys *et al.*, 1992).



These findings suggest that the P2Y purinoceptor represents a novel target for the development of oral antidiabetic drugs.

## 5.2. Cancer

Based on the potent growth inhibition of a variety of human and murine tumour cells by extracellular ATP, the administration of AMP or ATP to tumour-bearing murine hosts was shown to be associated with dramatic cytostatic and cytotoxic effects (Rapaport, 1993). Such anticancer activity is likely related to the activation of P2Z purinoceptors mediating cell killing through apoptosis (Pizzo *et al.*, 1993; Murgia *et al.*, 1992). The utilisation of ATP infusions against metastatic refractory cancers has already entered Phase I of human clinical trials. This clinical application is particularly intriguing since studies in cachectic tumour models have shown that the ATP-mediated tumour-killing activity is also associated with a variety of host-mediated anticancer activities, including significant inhibition of host weight loss (Rapaport, 1993).

## 5.3. Cystic Fibrosis

P2 purinoceptors have been shown to regulate ion transport in epithelial cells from a variety of different sources, including intestinal, lung and kidney epithelium, where ATP stimulates Cl<sup>-</sup> transport and alters Ca<sup>2+</sup> distribution (Burnstock, 1991). Cystic fibrosis (CF) is a lethal genetic disease characterised by defective regulation of chloride conductance in airway epithelia, which leads to the formation of underhydrated mucus obstructing the airways of patients with this disease (Noone and Knowles, 1993). Both ATP and UTP were found to be effective *in vitro* Cl<sup>-</sup> secretagogues in the nasal mucosa of human subjects, an effect that seemed secondary to the activation of phospholipase C and Ca<sup>2+</sup>-dependent phosphorylation of Cl<sup>-</sup> channels (Stuitts *et al.*, 1992). Most interestingly, regulation of Cl<sup>-</sup> conductance by ATP and UTP was preserved in CF nasal epithelia (Clarke and Boucher, 1992). On the basis of these results, it is intriguing to speculate that agonists at this P2-purinoceptor subtype (a P2Y<sub>2</sub> receptor?) might be of enormous clinical interest in normalising electrolyte and water secretion across CF airway epithelia.

## 5.4. Pulmonary Hypertension

In the lung, the most interesting target for drug development is represented by the vascular pulmonary P2 purinoceptor mediating ATP-induced vasoconstriction (McCormack *et al.*, 1993). This receptor seems different from the typical P2X purinoceptor responsible for excitatory (vasoconstrictor) responses, being equally activated by ATP and UTP (Rubino and Burnstock, 1994). The demonstration of a new purinoceptor involved in the control of pulmonary circulation is particularly intriguing, especially in view of the fact that attempts to lower pulmonary vascular resistance with pharmacological agents have been largely unsuccessful (McCormack *et al.*, 1993). Based on this, antagonists at the ATP/UTP vasoconstrictor lung purinoceptor might have therapeutic potential in hypoxic pulmonary vasoconstriction and in pulmonary hypertension.

## 5.5. Surfactant and Mucin Secretion

ATP also affects additional lung functional parameters. ATP has been demonstrated to increase mucin release from cultured airway goblet cells through the production of inositol phosphate (Kim *et al.*, 1993), suggesting a role for P2Y purinoceptors in maintaining lung visco-elastic properties and, consequently, in the defence against airborne particles. Phospholipase C-linked P2 purinoceptors also have been demonstrated to be expressed by Type II pneumocytes, the lung cell type responsible for surfactant secretion (Grise *et al.*, 1993). Activation of this receptor leads to increases of surfactant secretion into the bronchial lumen. Based on the hypothesis that, in some forms of pneumonia (Lachmann and Commers, 1993), the decreased respiratory performance can be attributed to deficiency of surfactant secretion, it would be possible to propose the use of ATP analogs to optimise surfactant secretion in these pathologies.

## 5.6. Renal Failure

Acute renal failure is commonly provoked by renal ischaemia. Current therapeutic strategies to improve renal blood flow are limited to resuscitation of the systemic circulation and infusion of low doses of dopamine. Failure of this line of management frequently leaves the patient requiring haemodialysis. P2X purinoceptors on the vascular smooth muscle of pre-glomerular (afferent) arterioles can represent an obvious target for antagonists that could enhance renal blood flow and help restore normal renal function. Likewise, analogues of ATP that stimulate P2Y purinoceptors could also increase renal blood flow (Churchill and Ellis, 1993a), although this may be opposed by P2Y purinoceptors mediating renin release (Churchill and Ellis, 1993b).

Renal infusion of ATP-MgCl<sub>2</sub> has been demonstrated to enhance post-ischaemic recovery of both glomerular and tubular functions (Siegel *et al.*, 1983; Hirasawa *et al.*, 1985), underlining the possible use of ATP analogues in preventing ischaemia-induced renal damage, although the mechanism of action is not fully understood. Beneficial effects of ATP-MgCl<sub>2</sub> have been demonstrated also against drug-induced nephrotoxicity (Sumpio *et al.*, 1985) and for kidney preservation before transplantation (Belzer *et al.*, 1983).

Interestingly, exogenous ATP stimulates the proliferation of mesangial cells in glomerulonephritis (Schulze-Lohoff *et al.*, 1992). Specific antagonists of P2 purinoceptors, therefore, may have a therapeutic potential in modulating the anti-inflammatory process of this serious disease.

## 5.7. Bone and Cartilage Diseases

ATP stimulates cartilage resorption through the activation of a P2 purinoceptor responsive to both ATP and UTP on chondrocytes (Caswell *et al.*, 1991). Interestingly, cytokines, which have been strongly implicated in the loss of cartilage extracellular matrix in rheumatoid arthritis and osteoarthritis, seem to act through enhancement of chondrocyte responsiveness to extracellular ATP (Leong *et al.*, 1993). Similarly, P2 purinoceptors have been demonstrated to be expressed also by osteoblasts (Kunagai *et al.*, 1991; Scholf *et al.*, 1992; Reimer and Dixon, 1992; Yu and Ferrer, 1993) and osteoclasts (Yu and Ferrer, 1993). In osteoblasts, P2 purinoceptor activation by both ATP and UTP has been associated with fast intracellular Ca<sup>2+</sup> pulses (a P2Y<sub>2</sub> receptor?). Since indirect stimulation of osteoclastic resorption is thought to be triggered by an initial elevation of [Ca<sup>2+</sup>]<sub>i</sub> in osteoblasts (Reimer and Dixon, 1992), it is intriguing to speculate that the blockade of the osteoblast receptor by selective P2Y-purinoceptor antagonists might be beneficial in pathologies characterised by excessive bone demineralisation such as osteoporosis.

## 5.8. Urinary Incontinence

ATP has been proposed to be the transmitter responsible for the atropine-resistant contractile response of isolated detrusor muscle elicited by transmural nerve stimulation (Burnstock *et al.*, 1972, 1978). This effect is exerted through the activation of P2X purinoceptors (Burnstock and Kennedy, 1985; Bo and Burnstock, 1990). In interstitial cystitis (a chronic bladder disorder characterised by incontinence), a pathological increase of detrusor muscle sensitivity to ATP analogs was demonstrated recently (Palcia *et al.*, 1993), opening interesting possibilities for selective P2X antagonists in the treatment of chronic bladder disorders characterised by dysuria.

## 5.9. Thrombosis

ATP has been reported to antagonise ADP-induced human platelet aggregation (Table 1; Cusack and Hourani, 1982), which might have interesting implications in the development of anti-thrombotic agents.

## 5.10. Ventricular Tachycardia

Both P<sub>1</sub>- and P<sub>2</sub>-purinoceptor antagonists have been recommended for paroxysmal ventricular tachycardia (Bellassen and Pelleg, 1984).

## 5.11. Gastro-Intestinal Tract

ATP has long been recognised as a nonadrenergic, noncholinergic inhibitory transmitter in the gastro-intestinal tract (Burnstock *et al.*, 1970) and more recently in sympathetic nerves supplying the intestine, where ATP is co-stored and co-released with noradrenaline (for a review, see Hoyle and Burnstock, 1991). P<sub>2</sub> purinoceptors (in most cases of the P<sub>2</sub>Y-subtype, mediating relaxation) so far have been identified in stomach and intestinal smooth muscle, in intestinal ganglia and in endothelial and smooth muscle cells of vessels supplying the gastro-intestinal tract (Hoyle and Burnstock, 1991; Burnstock *et al.*, 1994). In the next few years, these receptors are likely to be explored as possible targets for the development of new pharmacological agents with therapeutic potential in gastro-intestinal dysfunctions.

5.12. Centrally Targeted P<sub>2</sub> Purinoceptor Drugs

The recent demonstration of P<sub>2</sub> purinoceptors in the CNS (Table 5) also opens interesting opportunities for the development of novel pharmacological agents for the treatment of CNS disorders, such as epilepsy, depression and aging-associated neurodegenerative diseases (Burnstock, 1993; Williams, 1993).

## 5.13. Human Reproduction

The demonstration of ATP receptors on human spermatozoa (Forresta *et al.*, 1992) and amnion cells (Vander Kooy *et al.*, 1989) may suggest in the near future additional therapeutic implications for P<sub>2</sub>-purinoceptor agonists and antagonists in human reproduction.

## 6. FUTURE DEVELOPMENTS

Perhaps the most important area for future research, partly to substantiate our proposed P<sub>2</sub>-purinoceptor subclassification and also in relation to therapeutic development, is to identify selective antagonists for the different P<sub>2</sub>-purinoceptor subclasses defined in this review; antagonists that are effective *in vivo* are especially desirable.

*Acknowledgements*—The authors are grateful to Prof. E. Barnard, to Drs X. Bo, S. Cerni, C. H. V. Hoyle, K. A. Jacobson, C. Kennedy, P. Milner, V. Ralevic, A. L. Rubino, J. Simon, D. Williams and M. Williams, to Miss T. Webb for useful advice and suggestions, to Mrs A. Evans and C. Chiergato for typing the manuscript and to D. Christie for editorial help.

## NOTE ADDED IN PROOF

When this paper was already in press, two independent reports appeared in *Nature* on the cloning of two different ionotropic P<sub>2</sub>X-purinoceptors from rat *vas deferens* (Valera *et al.*, *Nature* 371: 516–519) and from PC12 cells (Brake *et al.*, *Nature* 371: 519–523). The first one is the receptor communicated by Suprenant in the present review. For both receptors, deduced membrane topology was strikingly different from that of other ionotropic transmitter receptors such as the acetylcholine nicotinic receptor. In both cases, deduced amino acid sequence contained two hydrophobic putative transmembrane domains with an intervening hydrophilic cysteine-rich loop of 270–287 amino acids, suggesting that P<sub>2</sub>X-purinoceptors may belong to an entirely different class of transmembrane signalling proteins. Consistently with our proposal on P<sub>2</sub>X-purinoceptor heterogeneity, the two receptors revealed different pharmacological profiles when expressed in oocytes and their transcripts showed different tissue distribution as revealed by *in situ* hybridisation analysis.

## REFERENCES

- Abbracchio, M. P., Cattabeni, F., Fredholm, B. B. and Williams, M. (1993) Purinoceptor nomenclature: a status report. *Drug Dev. Res.* 28: 207–213.  
 Allen, T. G. J. and Burnstock, G. (1990) The actions of adenosine 5'-triphosphate on guinea pig intracardiac neurons in culture. *Br. J. Pharmacol.* 100: 269–276.  
 Arthammar, P., Hallberg, A., Kindmark, H., Nilsson, T., Roisman, P. and Berggren, P. O. (1990) Extracellular ATP increases cytoplasmic free Ca<sup>2+</sup> concentration in clonal insulin-producing RINm5F cells. *Biochem. J.* 265: 203–211.  
 Bailey, S. J. and Hourani, S. M. O. (1992) Effects of purines on the longitudinal muscle of the rat colon. *Br. J. Pharmacol.* 105: 885–892.

- Barnard, E. A., Burnstock, G. and Webb, T. E. (1994) The G-protein coupled receptors for ATP and other nucleotides: a new receptor family. *Trends Pharmacol. Sci.* 15: 67–70.  
 Bean, B. P. (1990) ATP-activated channels in rat and bullfrog sensory neurons: concentration dependence and kinetics. *J. Neurosci.* 10: 1–10.  
 Bean, B. P. (1992) Pharmacology and electrophysiology of ATP-activated ion channels. *Trends Pharmacol. Sci.* 13: 87–90.  
 Belhassen, B. and Pelleg, A. (1984) Acute management of paroxysmal supraventricular tachycardia: verapamil, adenosine triphosphate or adenosine? *Am. J. Cardiol.* 54: 225–227.  
 Belzer, F. O., Sollinger, H. W., Glass, N. R., Miller, D. T., Hoffmann, R. M. and Southard, J. H. (1983) Beneficial effects of adenosine and phosphate in kidney preservation. *Transplantation* 36: 633–635.  
 Benham, C. D. and Tsien, R. W. (1987) A novel receptor-operated Ca<sup>2+</sup>-permeable channel activated by ATP in smooth muscle. *Nature* 328: 275–278.  
 Berne, R. M. (1963) Cardiac nucleotides in hypoxia: possible role in regulation of coronary blood flow. *Am. J. Physiol.* 204: 317–322.  
 Berridge, C. P., Hawkins, P. T., Stephens, L. R., Harden, T. K. and Downes, C. P. (1989) Phosphatidylinositol 4,5-bisphosphate hydrolysis in turkey erythrocytes is regulated by P<sub>2</sub>Y purinoceptors. *Molec. Pharmacol.* 35: 526–532.  
 Bertrand, G., Chapal, J., Loubatieres-Mariani, M. M. and Roye, M. (1987) Evidence for two different P<sub>2</sub> purinoceptors on  $\beta$  cell and pancreatic vascular bed. *Br. J. Pharmacol.* 91: 783–787.  
 Bjornsson, O. G., Monck, J. R. and Williamson, J. R. (1989) Identification of P<sub>2</sub>Y purinoceptors associated with voltage-activated cation channels in cardiac ventricular myocytes of the rat. *Eur. J. Biochem.* 186: 395–404.  
 Bo, X. and Burnstock, G. (1990) The effects of Bay K 8644 and nifedipine on the responses of rat urinary bladder to electrical field stimulation,  $\beta$ - $\gamma$ -methylene ATP and acetylcholine. *Br. J. Pharmacol.* 101: 494–498.  
 Bo, X., Fischer, B., Maillard, M., Jacobson, K. A. and Burnstock, G. (1994) Comparative studies on the affinities of ATP derivatives to P<sub>2</sub>Y-purinoceptors in rat urinary bladder. *Br. J. Pharmacol.* 112: 1151–1159.  
 Boyer, J. L., Cooper, C. L. and Harden, T. K. (1990) [<sup>3</sup>H]3-O-(4-Benzoyl) benzoyl ATP as a photoaffinity label for a phospholipase C-coupled P<sub>2</sub>Y-purinoceptor. *J. Biol. Chem.* 265: 13515–13520.  
 Bizzolara, A. L. and Burnstock, G. (1991) Endothelium-dependent and endothelium-independent vasodilatation of the hepatic artery of the rabbit. *Br. J. Pharmacol.* 103: 1206–1212.  
 Brown, H. A., Lazarewicz, E. R., Boucher, R. C. and Harden, T. K. (1991) Evidence that UTP and ATP regulate phospholipase C through a common extracellular 5'-nucleotide receptor in human airway epithelial cells. *Molec. Pharmacol.* 40: 648–655.  
 Bruner, G. and Murphy, S. (1990) ATP-evoked arachidonic acid mobilization in astrocytes is via a P<sub>2</sub>Y-purinoceptor. *J. Neurochem.* 55: 1569–1575.  
 Burnstock, G. (1972) Purinoceptor nerves. *Pharmacol. Rev.* 24: 509–581.  
 Burnstock, G. (1976) Do some nerve cells release more than one transmitter? *Neuroscience* 1: 239–248.  
 Burnstock, G. (1978) A basis for distinguishing two types of purinoceptor. In: *Cell Membrane Receptors for Drugs and Hormones: A Multidisciplinary Approach*, pp. 107–118, Straub, R. W. and Bolis, L. (eds) Raven Press, New York.  
 Burnstock, G. (1990) Co-transmission. The Fifth Heymans Memorial Lecture. *Arch. int. Pharmacodyn. Ther.* 304: 7–33.  
 Burnstock, G. (1991) Distribution and roles of purinoceptor subtypes. *Nucleosides Nucleotides* 10: 917–930.  
 Burnstock, G. (1993) Physiological and pathological roles of purines: an update. *Drug Dev. Res.* 28: 195–206.  
 Burnstock, G. and Kennedy, C. (1985) Is there a basis for distinguishing two types of P<sub>2</sub>Y-purinoceptor? *Gen. Pharmacol.* 16: 433–440.  
 Burnstock, G., Campbell, G., Satchell, D. G. and Smythe, A. (1970) Evidence that adenosine triphosphate or a related nucleotide is the transmitter substance released by non-adrenergic inhibitory nerves in the gut. *Br. J. Pharmacol.* 40: 668–688.  
 Burnstock, G., Dunsday, B. and Smythe, A. (1972) Atropine resistant excitation of the urinary bladder: the possibility of transmission via nerves releasing a purine nucleotide. *Br. J. Pharmacol.* 44: 451–456.  
 Burnstock, G., Cocks, T., Crowe, R. and Kasakov, L. (1978) Purinoceptor innervation of the guinea-pig urinary bladder. *Br. J. Pharmacol.* 63: 125–138.  
 Burnstock, G., Cusack, N. J., Mackenzie, I. and Meghji, P. (1983) Studies on the stereoselectivity of the P<sub>2</sub>Y-purinoceptor. *Br. J. Pharmacol.* 79: 907–913.  
 Burnstock, G., Cusack, N. J., Hills, J. M. and Meldrum, L. A. (1984) Effects of phosphorothioate analogues of ATP, ADP and AMP on guinea-pig taenia coli and urinary bladder. *Br. J. Pharmacol.* 82: 369–374.  
 Burnstock, G., Fischer, B., Hoyle, C. H. V., Maillard, M., Ziganshin, A. U., Bizzolara, A. L., von Isakowicz, A., Boyer, J. L., Kendall-Harden, T. and Jacobson, K. A. (1994) Structure activity relationships for derivatives of adenosine-5'-triphosphate as agonists at P<sub>2</sub> purinoceptors: heterogeneity within P<sub>2</sub>X and P<sub>2</sub>Y subtypes. *Drug Dev. Res.* 31: 206–219.  
 Busse, R., Ogilvie, A. and Pohl, U. (1988) Vasomotor activity of diadenosine triphosphate and diadenosine tetraphosphate in isolated arteries. *Am. J. Physiol.* 254: H828–832.  
 Carter, T. D., Hallam, T. J., Cusack, N. J. and Pearson, J. D. (1988) Regulation of P<sub>2</sub>Y-purinoceptor-mediated prostacyclin release from human endothelial cells by cytoplasmic calcium concentration. *Br. J. Pharmacol.* 95: 1181–1190.



- Castro, E., Pinior, J. and Miras-Portugal, M. T. (1992)  $\text{Ca}^{2+}$ -stores mobilization by diadenosine tetraphosphate, Ap4A, through a putative  $\text{P}_2\text{Y}$  purinoceptor in adrenal chromaffin cells. *Br. J. Pharmacol.* 106: 833-837.
- Carew, A. M., Leong, W. S. and Russell, R. G. G. (1991) Evidence for the presence of  $\text{P}_2$ -purinoceptors at the surface of human articular chondrocytes in monolayer culture. *Biochim. Biophys. Acta* 1074: 151-158.
- Churchill, P. C. and Ellis, V. R. (1993a) Pharmacological characterization of the renovascular  $\text{P}_2$  purinergic receptors. *J. Pharmacol. exp. Ther.* 265: 334-338.
- Churchill, P. C. and Ellis, V. R. (1993b) Purinergic  $\text{P}_2$  receptors stimulate renin secretion by rat renal cortical slices. *J. Pharmacol. exp. Ther.* 266: 160-163.
- Clarke, L. L. and Boucher, R. C. (1992) Chloride secretory response to extracellular ATP in human normal and cystic fibrosis nasal epithelia. *Am. J. Physiol.* 263 (Cell Physiol. 32): C348-C356.
- Cockcroft, S. and Stutchfield, J. (1989) The receptor for ATP and MethylLeuPhe are independently coupled to phospholipases C and  $\text{A}_2$  via G-protein(s). *Biochem. J.* 263: 715-723.
- Cooper, C. L., Morris, A. J. and Harden, T. K. (1989) Guanine nucleotide-sensitive interaction of a radiolabeled agonist with a phospholipase C-linked  $\text{P}_2\text{Y}$ -purinergic receptor. *J. Biol. Chem.* 264: 6202-6206.
- Comfield, L. J., Hu, S., Hurt, S. D. and Sills, M. A. (1992) [ $^3\text{H}$ ]-2-Phenylaminoadenosine ([ $^3\text{H}$ ]CV 1808) labels a novel adenosine receptor in rat brain. *J. Pharmacol. exp. Ther.* 263: 552-561.
- Cor, L. and Burnstock, G. (1991) Vasodilator response of coronary smooth muscle to the sympathetic co-transmitters noradrenaline and adenosine 5'-triphosphate. *Br. J. Pharmacol.* 104: 337-342.
- Cusack, N. J. (1993)  $\text{P}_2$  receptor: subclassification and structure-activity relationships. *Drug Dev. Res.* 28: 244-252.
- Cusack, N. J. and Hourani, S. M. O. (1982) Competitive inhibition by adenosine 5'-triphosphate of the actions on human platelets of 2-chloroadenosine 5'-diphosphate, 2-azido-adenosine 5'-diphosphate and 2-methylthioadenosine 5'-diphosphate. *Br. J. Pharmacol.* 77: 329-333.
- Cusack, N. J. and Hourani, S. M. O. (1990) Structure-activity relationships for adenosine nucleotide receptors on mast cells, human platelets and smooth muscle. In: *Purines in Cellular Signalling: Targets for New Drugs*, pp. 254-259. Jacobson, K. A., Daly, J. W. and Manganiello, V. (eds) Springer, New York.
- Cusack, N. J., Hickman, M. E. and Born, G. V. R. (1979) Effects of D- and L-enantiomers of adenosine, AMP and ADP and their 2-chloro and 2-azido-analogues on human platelets. *Proc. R. Soc. Lond. B* 206: 139-144.
- Cusack, N. J., Hourani, S. M. O., Loizou, G. D. and Welford, L. A. (1987) Pharmacological effects of isopolar phosphonate analogues of ATP on  $\text{P}_2$ -purinoceptors in guinea-pig, *laetia coli* and urinary bladder. *Br. J. Pharmacol.* 90: 791-795.
- Daly, J. W. (1982) Adenosine receptors: targets for future drugs. *J. med. Chem.* 25: 197-207.
- Daly, J. W., Nehlig, A. and Nicols, F. (1991) Physiological and pharmacological properties of adenosine: therapeutic implications. *Life Sci.* 49: 1435-1453.
- Davidson, J. S., Wakefield, I. K., Sohnius, U., Van der Merwe, P. A. and Millar, R. P. (1990) A novel extracellular nucleotide receptor coupled to phosphoinositide-C-inositol cells. *Endocrinology* 126: 80-87.
- Druy, A. N. and Szent-Gyorgyi, A. (1929) The physiological activity of adenine compounds with special reference to their action upon the mammalian heart. *J. Physiol. (Lond.)* 68: 213-237.
- Dulon, D., Moatéz, R. and Mollard, P. (1993) Characterization of  $\text{Ca}^{2+}$  signals generated by extracellular nucleotides in supporting cells of the Organ of Corti. *Cell Calcium* 14: 245-254.
- Edwards, J. C. (1964) *The Physiology of Synapses*. Springer-Verlag, Berlin.
- Edwards, F. A., Alsdair, J. G. and Colquhoun, D. (1992) ATP receptor-mediated synaptic currents in the central nervous system. *Nature* 359: 144-146.
- Elferink, J. G. R., de Koster, B. M., Boonen, G. J. J. C. and de Prieester, W. (1992) Inhibition of neurophil chemotaxis by purinoceptor agonists. *Arch. int. Pharmacodyn.* 317: 93-106.
- Fieber, L. A. and Adams, D. J. (1991) Adenosine triphosphate-evoked currents in cultured neurones dissociated from rat parasympathetic cardiac ganglia. *J. Physiol. (Lond.)* 434: 239-256.
- Fine, J., Cole, P. and Davidson, J. S. (1989) Extracellular nucleotides stimulate receptor calcium mobilization and inositol phosphate production in human fibroblasts. *Biochem. J.* 263: 371-376.
- Fink, J. S., Weaver, D. R., Rivekes, S. A., Peterfreund, R. A., Pollack, A. E., Adler, E. M. and Reppert, S. M. (1992) Molecular cloning of the rat  $\text{A}_1$  adenosine receptor: selective co-expression with  $\text{D}_1$  dopamine receptors in rat striatum. *Brain Res. molec. Brain Res.* 14: 186-195.
- Fischer, B., Boyer, J. L., Hoyle, C. H. V., Ziganishin, A. U., Brizzolara, A. L., Knight, G. E., Zimmel, J., Burnstock, G., Kendall-Harden, T. and Jacobson, K. A. (1993) Identification of potent  $\text{P}_2\text{Y}$  purinoceptor agonists: structure-activity relationships for 2-thioether derivatives of adenosine-5-triphosphate. *J. med. Chem.* 36: 3937-3946.
- Flodgaard, H. and Klenow, H. (1992) Abundant amounts of diadenosine 5',5''-p1,p4-tetraphosphate are present and releasable, but metabolically inactive in human platelets. *Biochem. J.* 208: 737-742.
- Foresta, C., Rossato, M. and Di Virgilio, F. (1992) Extracellular ATP is a trigger for the acrosome reaction in human spermatozoa. *J. Biol. Chem.* 267: 19443-19447.
- Fournier, F., Honoré, E., Collin, T. and Guilhaud, P. (1990) Ins(1,4,5)P3 formation and fluctuating chloride current response induced by external ATP in *Xenopus* oocytes injected with embryonic guinea-pig brain mRNA. *FEBS Lett.* 277: 205-208.
- Fredholm, B. B., Abbracchio, M. P., Burnstock, G., Daly, J. W., Harden, T. K., Jacobson, K. A. and Williams, M. (1994) Nomenclature and classification of purinoceptors: a report from the IUPHAR subcommittee. *Pharmac. Rev.* 46: 143-156.

- Frelin, C., Breitmayer, J. P. and Vigne, P. (1993) ADP induces inositol phosphate-independent intracellular  $\text{Ca}^{2+}$  mobilization in brain capillary endothelial cells. *J. Biol. Chem.* 268: 8787-8792.
- Gil-Rodrigo, C. E., Galdiz, B., Gandarias, J. M., Gomez, R. and Ainz, L. F. (1990) Characterization of the effects of adenosine, adenosine 5'-triphosphate and related purines on acid secretion in isolated rabbit gastric glands. *Pharmac. Res.* 2: 103-113.
- Gillespie, J. H. (1993) The biological significance of the linkages in adenosine triphosphoric acid. *J. Physiol. (Lond.)* 465: 345-349.
- Gilman, A. (1987) G-proteins: transducers of receptor-generated signals. *Annu. Rev. Biochem.* 56: 615-649.
- Goedcke, S. and Schrader, J. (1994) Cloning and expression analysis of the rat ATP receptor. *Drug Dev. Res.* 31: 273.
- Goldman, S. J., Gordon, E. L. and Slakey, L. L. (1986) Hydrolysis of diadenosine 5'-5''-p1,p4-triphosphate (Ap4A) by porcine aortic endothelial cells. *Circ. Res.* 59: 362-366.
- Gordon, J. L. (1986) Extracellular ATP: effects, sources and fate. *Biochem. J.* 233: 309-319.
- Green, H. N. and Stoner, H. B. (1990) *Biological Actions of the Adenosine Nucleotides*. Lewis, London.
- Grise, M., Gobran, L. J. and Roomey, S. A. (1993) Signal-transduction mechanisms of ATP-stimulated phosphatidylcholine secretion in rat type II pneumocytes: interactions between ATP and other surfactant secretagogues. *Biochim. Biophys. Acta* 1167: 85-93.
- Hausinger, D., Stiehle, T., Gerok, W., Tran-Thi, T. A. and Decker, K. (1987) Hepatocyte heterogeneity in response to extracellular ATP. *Eur. J. Biochem.* 169: 645-650.
- Hickman, S. E., Semnad, C. E., Field, M. and Silverstein, S. C. (1993) Expression of macrophage ATP and UTP receptors. *FASEB J.* 7: A712.
- Hilderman, R. H., Martin, M., Zimmerman, J. K. and Pivorum, E. B. (1991) Identification of a unique membrane receptor for adenosine 5',5''-p1,p4-tetraphosphate. *J. Biol. Chem.* 266: 6915-6918.
- Hillaire-Buys, D., Gross, R., Chapel, J., Ribes, G. and Loubatieres-Mariani, M. M. (1992)  $\text{P}_2\text{Y}$  purinoceptor responses of  $\beta$  cells and vascular bed are preserved in diabetic rat pancreas. *Br. J. Pharmacol.* 106: 610-615.
- Hillaire-Buys, D., Bertrand, G., Chapel, J., Puech, R., Ribes, G. and Loubatieres-Mariani, M. M. (1993) Stimulation of insulin secretion and improvement of glucose tolerance in rat and dog by the  $\text{P}_2\text{Y}$ -purinoceptor agonist, adenosine-5'-O-(2-thiodiphosphate). *Br. J. Pharmacol.* 109: 183-187.
- Hiraawa, H., Soeda, K., Ohake, Y., Oda, S., Kobayashi, S., Odaka, M. and Sato, H. (1985) Effects of ATP-MgCl<sub>2</sub> and ATP-Na<sub>2</sub> administration on renal function and cellular metabolism following renal ischemia. *Circ. Shock* 16: 337-346.
- Holton, F. A. and Holton, P. (1954) The capillary dilator substances in dry powders of spinal roots: a possible role of ATP in chemical transmission from nerve endings. *J. Physiol. (Lond.)* 126: 124-140.
- Holton, P. (1959) The liberation of adenosine triphosphate on antidromic stimulation of sensory nerves. *J. Physiol. (Lond.)* 145: 494-504.
- Honoré, E., Fournier, F., Collin, T., Nargeot, J. and Guilhaud, P. (1991) Functional expression of  $\text{P}_2\text{Y}$  purinoceptor in *Xenopus* oocyte injected with brain mRNA. *Pflügers Arch.* 418: 447-452.
- Hourani, S. M. O., Welford, L. A., Loizou, G. D. and Cusack, N. J. (1988) Adenosine 5'-[2-fluorodiphosphate] is a selective agonist at  $\text{P}_2$ -purinoceptors mediating relaxation of smooth muscle. *Eur. J. Pharmacol.* 147: 131-136.
- Hourani, S. M. O., Johnson, C. R. and Bailey, S. J. (1993) Desensitization of the  $\text{P}_2$ -purinoceptors on the rat colon muscularis mucosae. *Br. J. Pharmacol.* 110: 501-505.
- Houston, D. A., Burnstock, G. and Vanhoutte, P. M. (1987) Different  $\text{P}_2$ -purinergic receptor subtypes of endothelium and smooth muscle in canine blood vessels. *J. Pharmacol. exp. Ther.* 241: 501-506.
- Hoyle, C. H. V. (1990) Pharmacological activity of adenine dinucleotides in the periphery: possible receptor classes and transmitter function. *Gen. Pharmacol.* 21: 827-831.
- Hoyle, C. H. V. and Burnstock, G. (1991) ATP receptors and their physiological roles. In: *Adenosine in the Nervous System*, pp. 43-76. Stone, T. W. (ed.) Academic Press Ltd, London.
- Hoyle, C. H. V., Chapple, C. and Burnstock, G. (1989) Isolated human bladder: evidence for an adenine dinucleotide acting on  $\text{P}_2\text{Y}$ -purinoceptors and for purinergic transmission. *Eur. J. Pharmacol.* 174: 115-118.
- Hoyle, C. H., Knight, C. E. and Burnstock, G. (1990) Suramin antagonises responses to  $\text{P}_2$ -purinoceptor agonists and purinergic nerve stimulation in the guinea-pig urinary bladder and *laetia coli*. *Br. J. Pharmacol.* 99: 617-621.
- Illes, P. and Nörenberg, W. (1993) Neuronal ATP receptors and their mechanism of action. *Trends Pharmacol. Sci.* 14: 50-54.
- Inoue, K. and Nakazawa, K. (1992) ATP-receptor operated  $\text{Ca}^{2+}$  influx and catecholamine release from neuronal cells. *News Physiol. Sci.* 7: 56-58.
- Jacobson, K. A. (1990) Adenosine ( $\text{P}_1$ ) and ATP ( $\text{P}_2$ )-purinoceptors. In: *Comprehensive Medicinal Chemistry: The Rational Design, Mechanistic Study and Therapeutic Application of Chemical Compounds*, Vol. 3, *Membranes and Receptors*, pp. 601-642. Emmet, J. C. (ed.) Pergamon Press, Oxford.
- Jacobson, K. A., van Galen, P. J. M. and Williams, M. (1992) Adenosine receptors: pharmacology, structure-activity relationships, and therapeutic potential. *J. med. Chem.* 35: 407-422.
- Kaszkov, L. and Burnstock, G. (1983) The use of the slowly degradable analog,  $\alpha,\beta$ -methyleneATP to produce desensitisation of the  $\text{P}_2$ -purinoceptor: effect on non-adrenergic, non-cholinergic responses of the guinea-pig urinary bladder. *Eur. J. Pharmacol.* 86: 291-294.

- Kastritis, C. H. C., Salm, A. K. and McCarthy, K. (1992) Stimulation of the P<sub>2</sub> purinergic receptor on type I astroglia results in inositol phosphate formation and calcium mobilization. *J. Neurochem.* 58: 1277-1284.
- Keprens, S. (1993) The complex interaction of ATP and UTP with isolated hepatocytes. How many receptors? *Gen. Pharmac.* 24: 283-289.
- Keprens, S., Vandekerckhove, A. and De Wulf, H. (1992) Extracellular ATP and UTP exert similar effects on rat isolated hepatocytes. *Br. J. Pharmac.* 105: 475-479.
- Kerr, D. I. B. and Kranis, A. (1979) A new class of ATP antagonist. *Proc. Australian Physiol. Pharmac. Soc.* 10: 156P.
- Kim, K. C., Zheng, Q.-X. and Van Seuingen, I. (1993) Involvement of a signal transduction mechanism in ATP-induced mucin release from cultured airway goblet cells. *Am. J. Respir. Cell. molec. Biol.* 8: 121-125.
- Kidlin, K. I., Hisayama, T. and Takayama, I. (1992) Comparison of contractile mechanisms by carbachol and ATP in detrusor strips of rabbit urinary bladder. *Jpn. J. Pharmac.* 58: 219-229.
- Kishal, O. A., Marchenko, S. M. and Pidoplichko, V. I. (1983) Receptor for ATP in the membrane of mammalian sensory neurons. *Neurosci. Lett.* 35: 41-45.
- Kunagai, H., Sacktor, B. and Filburn, C. R. (1991) Purinergic regulation of cytosolic calcium and phosphoinositide metabolism in rat osteoblast-like osteosarcoma cells. *J. Bone Min. Res.* 6: 697-701.
- Lachmann, B. and Gommers, D. (1993) Is it rational to treat pneumonia with exogenous surfactant? *Eur. Respir. J.* 6: 1427-1428.
- Lambrecht, G., Friebe, T., Grimm, U., Windscheif, U., Bungardt, E., Hildebrandt, C., Baumert, H. G., Spatz-Kumbel, G. and Mutschler, E. (1992) PPADS, a novel functionally selective antagonist of P<sub>2</sub> purinoceptor-mediated responses. *Eur. J. Pharmac.* 217: 217-219.
- Langosch, J. M., Gebicke-Haerter, P. J., Norenberg, W. and Illies, P. (1994) Two classes of P2Y<sub>2</sub>-purinoceptors mediate a dual effect of ATP in rat microglial cells. *Drug. Dev. Res.* 31: 280.
- Lazarowski, E. R. and Harden, T. K. (1994) Identification of a uridine nucleotide-selective G-protein-linked receptor that activates phospholipase C. *J. Biol. Chem.* 269: 11830-11836.
- Lazarowski, E., Mason, S., Parr, E. and Boucher, R. (1994) Functional expression of a human P<sub>2U</sub>-purinergic receptor in 1321NI cells. *Drug. Dev. Res.* 31: 289.
- Leff, P., Humphries, R. G., Crack, B. E., Pollard, C. E. and McKechinie, K. (1994) New ATP analogues: pharmacological tools and potential drugs. *Drug. Dev. Res.* 31: 290.
- Leong, W. S., Russel, R. G. G. and Caswell, A. M. (1993) Induction of enhanced responsiveness of human articular chondrocytes to extracellular ATP by tumour necrosis factor- $\alpha$ . *Clin. Sci.* 16: 569-575.
- Li, G., Milani, D., Dunne, M. J., Pralong, W.-F., Theler, J.-M., Petersen, O. H. and Wollheim, C. B. (1991) Extracellular ATP causes Ca<sup>2+</sup>-dependent and -independent insulin secretion in RINm5F cells. *J. Biol. Chem.* 266: 3449-3457.
- Libert, F., Parmentier, M., Lefort, A., Dinsart, C., Van Sande, J., Maenhaut, C., Simons, M.-J., Dumont, J. E. and Vassart, G. (1989) Selective amplification and cloning of four new members of the G protein-coupled receptor family. *Science* 244: 569-572.
- Libert, F., Schiffmann, S. N., Lefort, A., Parmentier, M., Gerard, C., Dumont, J. E., Vanderaeghe, J.-J. and Vassart, G. (1991) The orphan receptor cDNA RDC7 encodes an A<sub>1</sub> adenosine receptor. *EMBO J.* 10: 1677-1682.
- Linden, J., Tucker, A. L. and Lynch, K. R. (1991) Molecular cloning of adenosine A<sub>1</sub> and A<sub>2</sub> receptors. *Trends Pharmac. Sci.* 12: 326-328.
- Linden, J., Jacobson, M. A., Hutchins, C. and Williams, M. (1994) Adenosine receptors. In: *Handbook of Receptors and Channels, Vol. 1, G-Protein-Coupled Receptors*, pp. 29-44. Perouka, S. J. (ed.) CRC Press, Boca Raton.
- Londos, C., Cooper, D. M. F. and Woolf, J. (1980) Subclasses of external adenosine receptors. *Proc. natn. Acad. Sci. U.S.A.* 77: 2551-2554.
- Lustig, K. D., Shiao, A. K., Brake, A. J. and Julius, D. (1993) Expression cloning of an ATP receptor from mouse neuroblastoma cells. *Proc. natn. Acad. Sci. U.S.A.* 90: 5113-5117.
- Lubje, J. and Ogilvie, A. (1983) Presence of diadenosine 5'-5''-p1-p4-triphosphate (Ap<sub>4</sub>A) in human platelets. *Biochem. biophys. Res. Commun.* 115: 253-260.
- Maenhaut, C., van Sande, J., Libert, F., Abramowicz, M., Parmentier, M., Vanderaeghe, J. J., Dumont, J. E., Vassart, G. and Schiffmann, S. (1990) RDC8 codes for an adenosine A<sub>2</sub> receptor with physiological constitutive activity. *Biochem. biophys. Res. Commun.* 173: 1169-1178.
- Maguire, M. H. and Satchell, D. G. (1979) The contribution of adenosine to the inhibitory actions of adenine nucleotides on the guinea-pig taenia coli: studies with phosphate modified adenine nucleotide analogs and diprydamole. *J. Pharmac. exp. Ther.* 211: 626-631.
- Mahan, L. C., McVittie, L. D., Smyk-Randall, E. M., Nakata, H., Mounsa, F. J., Jr., Gerfen, C. R. and Sibley, D. R. (1991) Cloning and expression of an A<sub>1</sub> adenosine receptor from rat brain. *Molec. Pharmac.* 40: 1-7.
- Manzini, S., Hoyle, C. H. V. and Burnstock, G. (1986) An electrophysiological analysis of the effect of reactive blue 2, a putative P<sub>2</sub>-purinoceptor antagonist, on inhibitory junction potentials of rat caecum. *Eur. J. Pharmac.* 127: 197-204.
- Martin, G. N., Thom, S. A. M. and Sever, P. S. (1991) The effects of adenosine triphosphate (ATP) and related purines on human isolated subcutaneous and omental resistance arteries. *Br. J. Pharmac.* 102: 645-650.
- Martin, W., Cusack, N. J., Carleton, J. S. and Gordon, J. L. (1985) Specificity of P<sub>2</sub>-purinoceptor that mediates endothelium-dependent relaxation of the pig aorta. *Eur. J. Pharmac.* 108: 295-299.

- Mason, S. J., Paradiso, A. M. and Boucher, R. C. (1991) Regulation of transepithelial ion transport and intracellular calcium by extracellular ATP in human normal and cystic fibrosis airway epithelium. *Br. J. Pharmac.* 103: 1649-1656.
- McCormack, D. G., Crawley, D. E. and Evans, T. W. (1993) New perspectives in the pulmonary circulation and hypoxic pulmonary vasoconstriction. *Pulm. Pharmac.* 6: 97-108.
- Meyeroth, W., Muller-Brechlin, R. and Richter, D. (1991) Molecular cloning of a novel putative G-protein coupled receptor expressed during rat spermiogenesis. *FEBS Lett.* 284: 155-160.
- Murgia, M., Pizzo, P., Zanollo, P., Zamboni, A. and Di Virgilio, F. (1992) *In vitro* cytotoxic effects of extracellular ATP. *ATL4* 20: 66-70.
- Murgia, M., Haanu, S., Pizzo, P., Ripa, M. and Di Virgilio, F. (1993) Oxidized ATP—An irreversible inhibitor of the macrophage purinergic P<sub>2</sub> receptor. *J. Biol. Chem.* 268: 8199-8203.
- Murphy, P. M. and Tiffany, H. L. (1990) Characterization of phagocyte P<sub>2</sub> nucleotide receptors expressed in *Xenopus* oocytes. *J. Biol. Chem.* 265: 11615-11621.
- Murria, R. J. A. and Boarder, M. R. (1992) Neuronal 'nucleotide' receptor linked to phospholipase C and phospholipase D? Stimulation of PC12 cells by ATP analogues and UTP. *Molec. Pharmac.* 41: 561-568.
- Nakagawa, T., Akaike, N., Kimitsuki, T., Komura, S. and Arima, T. (1990) ATP-induced current in isolated outer hair cells of guinea pig cochlea. *J. Neurophysiol.* 5: 1068-1074.
- Noone, P. G. and Knowles, M. R. (1993) Trends in therapy of abnormal airway epithelial ion and liquid transport. *Monaldi Arch. Chest. Dis.* 48: 140-143.
- O'Connor, S. E., Wood, B. E. and Leff, P. (1990) Characterisation of P<sub>2X</sub>-receptors in rabbit isolated ear artery. *Br. J. Pharmac.* 101: 640-644.
- Ogilvie, A., Lubje, J., Pohl, U. and Busse, R. (1989) Identification and partial characterization of an adenosine (5')tetraphosphate (5') adenosine hydrolase on intact bovine aortic endothelial cells. *Biochem. J.* 259: 97-103.
- Okajima, F., Sato, K. and Kondo, Y. (1989) P<sub>2</sub>-purinergic agonists activate phospholipase C in a guanine nucleotide- and Ca<sup>2+</sup>-dependent manner in FRTL-5 thyroid cell membranes. *FEBS Lett.* 253: 132-136.
- Palen, S., Arribani, W., Ostardo, E., Tris, D. G. and Pietra, C. (1993) Evidence for purinergic neurotransmission in human urinary bladder affected by interstitial cystitis. *J. Urol.* 150: 2007-2012.
- Parr, C. E., Sullivan, D. M., Paradiso, A. M., Lazarowski, E. R., Burch, L. H., Olsen, J. C., Erb, L., Weisman, G. A., Boucher, R. C. and Turner, J. T. (1994) Cloning and expression of a human P2U nucleotide receptor, a target for cystic fibrosis pharmacotherapy. *Proc. natn. Acad. Sci. U.S.A.* 91: 3275-3279.
- Pavenstahl, H., Lindeman, V., Lindemann, S., Kunzelmann, S., Speth, M. and Greger, R. (1991) Effect of depolarizing and hyperpolarizing agents on the membrane potential difference of primary cultures of rabbit aorta vascular smooth muscle cells. *Eur. J. Physiol.* 419: 69-75.
- Pfeilschifter, J. (1990) Comparison of extracellular ATP and UTP signalling in rat renal mesangial cells—No indications for the involvement of separate purino- and pyrimidino-receptors. *Biochem. J.* 272: 469-472.
- Pintor, J. and Miras-Portugal, M. T. (1993) Diadenosine polyphosphates (Ap<sub>4</sub>A) as new neurotransmitters. *Drug Dev. Res.* 28: 259-262.
- Pintor, J., Torres, M., Castro, E. and Miras-Portugal, M. T. (1991a) Characterization of diadenosine tetraphosphate (Ap<sub>4</sub>A) binding sites in cultured chromaffin cells: evidence for a P<sub>2Y</sub> site. *Br. J. Pharmac.* 103: 1980-1984.
- Pintor, J., Torres, M. and Miras-Portugal, M. T. (1991b) Carbachol induced release of diadenosine polyphosphates—Ap<sub>4</sub>A and Ap<sub>3</sub>A—From perfused bovine adrenal medulla and isolated chromaffin cells. *Life Sci.* 48: 2317-2324.
- Pintor, J., Diaz-Rey, M. A. and Miras-Portugal, M. T. (1993) Ap<sub>4</sub>A and ADP- $\beta$ -S binding to P<sub>2</sub> purinoceptors present on rat brain synaptic terminals. *Br. J. Pharmac.* 108: 1094-1099.
- Pizzo, P., Murgia, M., Zamboni, A., Zanollo, P., Bronte, V., Pietrobon, D. and Di Virgilio, F. (1993) Role of P<sub>2</sub> purinergic receptors in ATP-mediated killing of tumor necrosis factor (TNF)-sensitive and TNF-resistant L929 fibroblasts. *J. Immun.* 109: 3372-3378.
- Purkiss, J. R., Wilkinson, G. F. and Boarder, M. R. (1993) Evidence for a nucleotide receptor on adrenal medullary endothelial cells linked to phospholipase C and phospholipase D. *Br. J. Pharmac.* 108: 1031-1037.
- Ralevic, V. and Burnstock, G. (1991) Effects of purines and pyrimidines on the rat mesenteric arterial bed. *Circ. Res.* 69: 1583-1590.
- Rapaport, E. (1993) Anticancer activities of adenine nucleotides in tumor bearing hosts. *Drug Dev. Res.* 28: 428-431.
- Rapaport, E. and Zamecnik, P. C. (1976) Presence of diadenosine 5'-5''-p1-p4-tetraphosphate (AP<sub>4</sub>A) in mammalian cells in levels varying widely with proliferative activity of the tissue: a possible positive pleiotropic activator. *Proc. natn. Acad. Sci. U.S.A.* 73: 3984-3988.
- Reed, J. K., de Souza, L., Raha, S. and Moore, H. (1994) ATP and UTP activate distinct signalling pathways in PC12 cells. *Drug Dev. Res.* 31: 312.
- Reimer, W. J. and Dixon, S. J. (1992) Extracellular nucleotides elevate [Ca<sup>2+</sup>]<sub>i</sub> in rat osteoblastic cells by interaction with two receptor subtypes. *Am. J. Physiol.* 263 (Cell Physiol. 32): C1040-C1048.
- Rubino, A. and Burnstock, G. (1994) A new purinergic receptor may contribute to the control of pulmonary circulation of the rat. *Drug Dev. Res.* 31: 315.
- Russell, S. N., Horner, M. A., Westfall, D. P., Buxton, I. L. O. and Horowitz, B. (1993) Characterization and functional expression of the ATP receptor in vas deferens smooth muscle. *Biophys. J.* 64: A84.



- Saig, B., Milton, D., Allain, H., Raul, B. and Van Den Driessche, J. (1990) Constriction of the smooth muscle of rat tail and femoral arteries and dog saphenous vein is induced by uridine triphosphate via pyrimidinopceptors, and by adenosine triphosphate via P<sub>2</sub> purinopceptors. *Blood Vessels* 27: 352-364.
- Saig, B., Milton, D., Shaoon, V., Allain, H., Raul, B. and Vandendor, J. (1992) Newly evidenced pyrimidinopceptors and the P<sub>2</sub> purinopceptors are present on the vascular smooth-muscle and respectively mediate the UTP-induced and ATP-induced contractions of the dog maxillary internal vein. *Res. Commun. CP* 76: 89-94.
- Salter, M. W. and Hicks, J. L. (1994) ATP-evoked increases in intracellular calcium in neurones and glia from the dorsal spinal cord. *J. Neurosci.* 14: 1563-1575.
- Salvatore, C. A., Luneau, C. J., Johnson, R. G. and Jacobson, M. A. (1992) Molecular cloning and characterization of human A<sub>1</sub> adenosine receptors. *Int. J. Purine Pyrim. Res.* 3: 82.
- Satchell, D. G. and Maguire M. H. (1975) Inhibitory effects of adenosine nucleotide analogs on the isolated guinea-pig taenia coli. *J. Pharmac. exp. Ther.* 195: 540-548.
- Schoff, C., Cuthbertson, K. S. R., Walsh, C. A., Mayne, C., Cobbold, P., Von Zur Muhlen, A., Hesch, R. D. and Gallagher, J. A. (1992) Evidence for P<sub>2</sub> purinopceptors on human osteoblast-like cells. *J. Bone Min. Res.* 5: 485-491.
- Schulze-Lohoff, E., Zanner, S., Ogilvie, A. and Stetzel, R. B. (1993) Extracellular ATP stimulates proliferation of cultured mesangial cells via P<sub>2</sub> purinergic receptors. *Am. J. Physiol.* 263: F374-F383.
- Shen, K.-Z. and North, R. A. (1993) Excitation of rat locus coeruleus neurons by adenosine 5'-triphosphate: ionic mechanism and receptor characterization. *J. Neurosci.* 13: 894-899.
- Siegel, N. J., Avison, M. J., Reilly, H. F., Alger, J. R. and Shulman, R. G. (1983) Enhanced recovery of renal ATP with postischemic infusion of ATP-MgCl<sub>2</sub> determined by <sup>31</sup>P-NMR. *Am. J. Physiol.* 245 (Renal Fluid Electrolyte Physiol. 14): F530-F534.
- Silinsky, E. M. (1989) Adenosine derivatives and neuronal function. *The Neurosciences* 1: 155-165.
- Stehle, J. H., Rivkees, S. A., Lee, J. J., Weaver, D. R., Deeds, J. D. and Reppert, S. M. (1992) Molecular cloning and expression of the cDNA for a novel A<sub>2</sub>-adenosine receptor subtype. *Molec. Endocrin.* 6: 384-393.
- Stone, T. (1992) Therapeutic potential of adenosine. *Script Medicae July/August*: 41-43.
- Stone, T. W. and Perkins, M. N. (1981) Adenosine dinucleotide effect on rat cortical neurones. *Brain Res.* 229: 241-245.
- Stuts, M. J., Chinet, T. C., Mason, S. J., Fullon, J. M., Clarke, L. L. and Boucher, R. C. (1992) Regulation of Cl<sup>-</sup> channels in normal and cystic fibrosis airway epithelial cells by extracellular ATP. *Proc. natn. Acad. Sci. U.S.A.* 89: 1621-1625.
- Sumplio, B. E., Chaudry, I. H. and Baue, A. E. (1985) Reduction of the drug induced nephrotoxicity by ATP-MgCl<sub>2</sub>. *J. Surg. Res.* 38: 429-437.
- Tawada, Y., Furukawa, K.-I. and Shigetawa, M. (1987) ATP-induced calcium transient in cultured rat aortic smooth muscle cells. *J. Biochem.* 102: 1499-1509.
- Theobald, R. J., Jr. (1992) Subclasses of purinopceptors in feline bladder. *Eur. J. Pharmac.* 229: 125-130.
- Thomas, S. A., Zawisa, M. J., Lin, X. and Hume, R. I. (1991) A receptor that is highly specific for extracellular ATP in developing chick skeletal muscle *in vitro*. *Br. J. Pharmac.* 103: 1963-1969.
- Tschöpl, M., Harris, L., Norenberg, W. and Illies, P. (1992) Excitatory effects of adenosine 5'-triphosphate on rat locus coeruleus neurones. *Eur. J. Pharmac.* 213: 71-77.
- Ueno, S., Harata, N., Inoue, K. and Akaike, N. (1992) ATP-gated current in dissociated rat nucleus solitarius neurons. *J. Neurophysiol.* 68: 778-786.
- van Calcar, D., Muller, M. and Hamprecht, B. (1979) Adenosine regulates via two different types of receptors, the accumulation of cyclic AMP in cultured brain cells. *J. Neurosci.* 33: 999-1005.
- Vander Kooy, D., Dubyak, G. R., Moore, R. M. and Moore, J. J. (1988) Adenosine triphosphate activates the phospholipase-C cascade system in human amnion cells without increasing prostaglandin production. *Endocrinology* 124: 2005-2012.
- von Kugelgen, I. and Starke, K. (1990) Evidence for two separate vasoconstriction-mediating nucleotide receptors, both distinct from the P<sub>2</sub>-receptor, in rabbit basilar artery: a receptor for pyrimidine nucleotides and a receptor for purine nucleotides. *Naunyn-Schmiedeberg's Arch. Pharmac.* 341: 538-546.
- von Kugelgen, I., Hausinger, D. and Starke, K. (1987) Evidence for a vasoconstriction-mediating receptor for UTP, distinct from the P<sub>2</sub> purinopceptor, in rabbit ear artery. *Naunyn-Schmiedeberg's Arch. Pharmac.* 336: 556-560.
- von Kugelgen, I., Bultmann, R. and Starke, K. (1990) Interaction of adenosine nucleotides, UTP and suramin in mouse vas deferens: suramin-sensitive and suramin-insensitive components in the contractile effect of ATP. *Naunyn-Schmiedeberg's Arch. Pharmac.* 342: 198-205.
- von Kugelgen, I., Späth, L. and Starke, K. (1994) Presynaptic P<sub>2</sub>-purinopceptors inhibiting the release of noradrenaline in rat brain cortex slices. *Drug Dev. Res.* 31: 332.
- Walker, B. A. M., Hagenlocker, B. E., Douglas, V. K., Tarapchak, S. J. and Ward, P. A. (1991) Nucleotide responses of human neutrophils. *Lab. Invest.* 64: 105-112.
- Webb, T. E., Simon, J., Krishak, B. J., Bateson, A. N., Smart, T. G., King, B. F., Burnstock, G. and Barnard, E. A. (1993) Cloning and functional expression of a brain G-protein-coupled ATP receptor. *FEBS Lett.* 324: 219-225.
- Westfall, D. P., Shinozuka, K., Forsyth, K. N. and Bjur, R. A. (1990) Presynaptic purine receptors. *Ann. N.Y. Acad. Sci.* 604: 130-135.
- Wilklund, N. P. and Gustafsson, L. E. (1988) Indications for P<sub>2</sub>-purinopceptors subtypes in guinea pig smooth muscle. *Eur. J. Pharmac.* 148: 361-370.
- Williams, M. (1987) Purine receptors in mammalian tissues: pharmacology and functional significance. *Ann. Rev. Pharmac. Toxicol.* 27: 315-345.
- Williams, M. (1993) Purinergic drugs: opportunities in the 1990s. *Drug Dev. Res.* 28: 438-444.
- Williams, M. (1994) Purinopceptor nomenclature: challenges for the future. In: *Adenosine and Adenine Nucleotides: From Molecular Biology to Integrative Physiology*, Belandini L. and Pelleg A. (eds) Kluwer, Norwell, MA: in press.
- Xing, M., Thevenod, F. and Mattern, R. (1992) Dual regulation of arachidonic acid release by P<sub>2</sub> purinergic receptors in dibutyryl cyclic AMP-differentiated HL60 cells. *J. Biol. Chem.* 267: 6602-6610.
- Yamada, M., Hamamori, Y., Akita, H. and Yokoyama, M. (1992) P<sub>2</sub>-purinopceptor activation stimulates phosphoinositide hydrolysis and inhibits accumulation of cAMP in cultured ventricular myocytes. *Circ. Res.* 70: 477-485.
- Yu, H. and Ferrer, J. (1993) ATP induces an intracellular calcium pulse in osteoclasts. *Biochem. biophys. Res. Commun.* 191: 357-363.
- Zamecnik, P. C., Byung, K., Gao, M. J., Taylor, G. and Blackburn, M. (1992) Analogues of diadenosine 5',5''-p<sub>1</sub>,p<sub>4</sub>-tetraphosphate (Ap<sub>4</sub>A) as potential anti-platelet-aggregation agents. *Proc. natn. Acad. Sci. U.S.A.* 89: 2370-2373.
- Zhou, F. Q.-Y., Olat, M. E., Li, C., Johnson, R. A., Stiles, G. L. and Civelli, O. (1992) Molecular cloning and characterization of a novel adenosine receptor: the A<sub>3</sub> adenosine receptor. *Proc. natn. Acad. Sci. U.S.A.* 89: 7432-7436.
- Ziganshin, A. U., Hoyle, C. H. V., Bo, X., Lambrecht, G., Mutschler, E., Baumann, H. G. and Burnstock, G. (1993) PPADS selectively antagonizes P<sub>2</sub>-purinopceptor-mediated responses in the rabbit urinary bladder. *Br. J. Pharmac.* 110: 1491-1495.
- Ziganshin, A. U., Hoyle, C. H. V. and Burnstock, G. (1994) Ecto-enzymes and metabolism of extracellular ATP. *Drug Dev. Res.* 32: 134-146.
- Ziganshin, A. U., Hoyle, C. H. V., Lambrecht, G., Baumann, H. G., Mutschler, E. and Burnstock, G. (1994b) Selective antagonism by PPADS at P<sub>2</sub>-purinopceptors in rabbit isolated blood vessels. *Br. J. Pharmac.* 111: 923-929.
- Zimmer, J., Jarlebark, L., van Galen, P. J. M., Jacobson, K. A. and Heilbronn, E. (1993) Synthesis and biological activity of novel 2-thio derivatives of ATP. *Nucleosides Nucleotides* 12: 1-20.